

# The Manin constant and $p$ -adic bounds on denominators of the Fourier coefficients of newforms at cusps

Abhishek Saha

(Joint work with K Česnavičius and M Neururer)

4th June, 2020

## Some facts on cusps for $\Gamma_0(N)$

- Any **cuspidal**  $\mathfrak{c} \in X_0(N)(\mathbb{C})$  is equivalent to

$$\mathfrak{c} = \frac{a}{L}, \text{ for some } L|N, \gcd(a, L) = 1.$$

We call  $L$  the *denominator* of  $\mathfrak{c}$ . There are exactly  $\phi(\gcd(L, N/L))$  cusps of denominator  $L$ .

## Some facts on cusps for $\Gamma_0(N)$

- Any **cuspidal**  $\mathfrak{c} \in X_0(N)(\mathbb{C})$  is equivalent to

$$\mathfrak{c} = \frac{a}{L}, \text{ for some } L|N, \gcd(a, L) = 1.$$

We call  $L$  the *denominator* of  $\mathfrak{c}$ . There are exactly  $\phi(\gcd(L, N/L))$  cusps of denominator  $L$ .

- The **width** of a cusp  $\mathfrak{c} = \frac{a}{L}$  equals

$$w(\mathfrak{c}) = \frac{N}{\gcd(L^2, N)}.$$

$w(\mathfrak{c})$  is the smallest integer  $w$  such that  $\begin{pmatrix} a & * \\ L & * \end{pmatrix} \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & * \\ L & * \end{pmatrix}^{-1} \in \Gamma_0(N)$ .

## Some facts on cusps for $\Gamma_0(N)$

- Any **cuspidal**  $\mathfrak{c} \in X_0(N)(\mathbb{C})$  is equivalent to

$$\mathfrak{c} = \frac{a}{L}, \text{ for some } L|N, \gcd(a, L) = 1.$$

We call  $L$  the *denominator* of  $\mathfrak{c}$ . There are exactly  $\phi(\gcd(L, N/L))$  cusps of denominator  $L$ .

- The **width** of a cusp  $\mathfrak{c} = \frac{a}{L}$  equals

$$w(\mathfrak{c}) = \frac{N}{\gcd(L^2, N)}.$$

$w(\mathfrak{c})$  is the smallest integer  $w$  such that  $\begin{pmatrix} a & * \\ L & * \end{pmatrix} \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & * \\ L & * \end{pmatrix}^{-1} \in \Gamma_0(N)$ .

- The **Atkin-Lehner involutions**: Let  $\mathfrak{c} = \frac{a}{L}$  be a cusp. Then there exists an Atkin-Lehner involution taking  $\mathfrak{c}$  to a cusp of denominator  $L'$  iff  $\text{val}_p(L') \in \{\text{val}_p(L), \text{val}_p(N) - \text{val}_p(L)\}$  for each  $p|N$ .

## The main question

- Let  $f = \sum_{n>0} a_f(n)q^n$ ,  $q = e^{2\pi iz}$  be a holomorphic **newform** of weight  $k$ , level  $N$ , trivial character.
- Normalize  $a_f(1) = 1$ . Then it well-known that all  $a_f(n) \in \overline{\mathbb{Z}}$ .

## The main question

- Let  $f = \sum_{n>0} a_f(n)q^n$ ,  $q = e^{2\pi iz}$  be a holomorphic **newform** of weight  $k$ , level  $N$ , trivial character.
- Normalize  $a_f(1) = 1$ . Then it well-known that all  $a_f(n) \in \overline{\mathbb{Z}}$ .
- Fourier expansion at  $\mathfrak{c}$ : Let  $\mathfrak{c} = \gamma\infty$  with  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ .

$$(f|_k\gamma)(z) = \sum_{n \geq 0} a_f(n; \mathfrak{c}) q^{\frac{n}{w(\mathfrak{c})}}.$$

**Note:**  $a_f(n; \mathfrak{c})$  only well-defined up to a  $w(\mathfrak{c})$ 'th root of unity.

- What can we say about the “denominators” of  $a_f(n; \mathfrak{c})$ ?

## The main question

- Let  $f = \sum_{n>0} a_f(n)q^n$ ,  $q = e^{2\pi iz}$  be a holomorphic **newform** of weight  $k$ , level  $N$ , trivial character.
- Normalize  $a_f(1) = 1$ . Then it well-known that all  $a_f(n) \in \overline{\mathbb{Z}}$ .
- Fourier expansion at  $\mathfrak{c}$ : Let  $\mathfrak{c} = \gamma\infty$  with  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ .

$$(f|_k\gamma)(z) = \sum_{n \geq 0} a_f(n; \mathfrak{c}) q^{\frac{n}{w(\mathfrak{c})}}.$$

**Note:**  $a_f(n; \mathfrak{c})$  only well-defined up to a  $w(\mathfrak{c})$ 'th root of unity.

- What can we say about the “denominators” of  $a_f(n; \mathfrak{c})$ ?

For a prime  $p$ , we are interested in good *lower* bounds for

$$\mathrm{val}_p(f|_{\mathfrak{c}}) := \inf_{n \geq 0} (\mathrm{val}_p(a_f(n; \mathfrak{c}))).$$

Here,  $\mathrm{val}_p: \overline{\mathbb{Q}}_p \rightarrow \mathbb{Q} \cup \{\infty\}$  is the  $p$ -adic valuation with  $\mathrm{val}_p(p) = 1$ , extended to  $\overline{\mathbb{C}}$  via any fixed choice of isomorphism  $\mathbb{C} \simeq \overline{\mathbb{Q}}_p$ .

Let  $f$  be a normalized newform for  $\Gamma_0(N)$  of weight  $k$ .

Find good *lower* bounds for  $\text{val}_p(f|_c) := \inf_{n \geq 0} (\text{val}_p(a_f(n; c)))$ .

- 1 Clearly,  $\text{val}_p(f|_\infty) = 0$ .

Let  $f$  be a normalized newform for  $\Gamma_0(N)$  of weight  $k$ .

Find good *lower* bounds for  $\text{val}_p(f|_c) := \inf_{n \geq 0} (\text{val}_p(a_f(n; c)))$ .

- 1 Clearly,  $\text{val}_p(f|_\infty) = 0$ .
- 2 **The  $q$ -expansion principle:** If the Fourier coefficients at infinity lie in a ring  $R$ , then the Fourier coefficients at any cusp lie in  $R[1/N, e^{\frac{2\pi i}{N}}]$ .  
In particular,  $\text{val}_p(f|_c) = 0$  if  $p \nmid N$ .

Let  $f$  be a normalized newform for  $\Gamma_0(N)$  of weight  $k$ .

Find good *lower* bounds for  $\text{val}_p(f|_c) := \inf_{n \geq 0} (\text{val}_p(a_f(n; c)))$ .

- 1 Clearly,  $\text{val}_p(f|_\infty) = 0$ .
- 2 **The  $q$ -expansion principle:** If the Fourier coefficients at infinity lie in a ring  $R$ , then the Fourier coefficients at any cusp lie in  $R[1/N, e^{\frac{2\pi i}{N}}]$ . In particular,  $\text{val}_p(f|_c) = 0$  if  $p \nmid N$ .
- 3 Suppose  $N$  is **squarefree** and  $p|N$ . Then using Atkin-Lehner operators, all cusps can be moved to  $\infty$ . An easy calculation now shows that:

$$\text{val}_p(f|_c) = \begin{cases} -\frac{k}{2} & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1. \end{cases}$$

Let  $f$  be a normalized newform for  $\Gamma_0(N)$  of weight  $k$ .

Find good *lower* bounds for  $\text{val}_p(f|_c) := \inf_{n \geq 0} (\text{val}_p(a_f(n; c)))$ .

- 1 Clearly,  $\text{val}_p(f|_\infty) = 0$ .
- 2 **The  $q$ -expansion principle:** If the Fourier coefficients at infinity lie in a ring  $R$ , then the Fourier coefficients at any cusp lie in  $R[1/N, e^{\frac{2\pi i}{N}}]$ . In particular,  $\text{val}_p(f|_c) = 0$  if  $p \nmid N$ .
- 3 Suppose  $N$  is **squarefree** and  $p|N$ . Then using Atkin-Lehner operators, all cusps can be moved to  $\infty$ . An easy calculation now shows that:

$$\text{val}_p(f|_c) = \begin{cases} -\frac{k}{2} & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1. \end{cases}$$

- 4 Nothing much previously known for general  $N$ . Some generic bounds exist due to Conrad using intersection theory on regular stacky surfaces, but are quite weak and have other issues.

Let  $f$  be a normalized newform for  $\Gamma_0(N)$  of weight  $k$ .

Find good *lower* bounds for  $\text{val}_p(f|_c) := \inf_{n \geq 0} (\text{val}_p(a_f(n; c)))$ .

- 1 Clearly,  $\text{val}_p(f|_\infty) = 0$ .
- 2 **The  $q$ -expansion principle:** If the Fourier coefficients at infinity lie in a ring  $R$ , then the Fourier coefficients at any cusp lie in  $R[1/N, e^{\frac{2\pi i}{N}}]$ . In particular,  $\text{val}_p(f|_c) = 0$  if  $p \nmid N$ .
- 3 Suppose  $N$  is **squarefree** and  $p|N$ . Then using Atkin-Lehner operators, all cusps can be moved to  $\infty$ . An easy calculation now shows that:

$$\text{val}_p(f|_c) = \begin{cases} -\frac{k}{2} & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1. \end{cases}$$

- 4 Nothing much previously known for general  $N$ . Some generic bounds exist due to Conrad using intersection theory on regular stacky surfaces, but are quite weak and have other issues.
- 5 For the general case, it suffices (thanks to AL operators) to restrict to cusps of denominator  $L$  such that  $L^2|N$ .

## Examples

$$N = 2^3 \cdot 3, k = 2, p = 2$$

$$f = q - q^2 + q^4 + q^5 + 2q^7 + \dots$$

$$f|_2\left(\frac{1}{2} \frac{1}{3}\right) = \frac{1}{6} \left( iq^{\frac{1}{6}} + iq^{\frac{1}{2}} - 2iq^{\frac{5}{6}} + \dots \right).$$

$$\text{So } \text{val}_2(f|_{1/2}) = -1.$$

## Examples

$$N = 2^3 \cdot 3, k = 2, p = 2$$

$$f = q - q^2 + q^4 + q^5 + 2q^7 + \dots$$

$$f|_2 \left( \frac{1}{2} \frac{1}{3} \right) = \frac{1}{6} \left( iq^{\frac{1}{6}} + iq^{\frac{1}{2}} - 2iq^{\frac{5}{6}} + \dots \right).$$

$$\text{So } \text{val}_2(f|_{1/2}) = -1.$$

$$N = 2 \cdot 3^5, k = 2, p = 3$$

$$f = q - q^2 + q^4 + 3q^5 - 4q^7 + \dots$$

$$f|_2 \left( \frac{1}{3} \frac{-1}{-2} \right) = \frac{1}{54} \left( \zeta_{162}^{25} q^{\frac{1}{54}} + \zeta_{162}^{50} q^{\frac{2}{54}} + \zeta_{162}^{19} q^{\frac{4}{54}} + \dots \right)$$

$$f|_2 \left( \frac{1}{9} \frac{1}{10} \right) = \frac{1}{6} \left( \zeta_{54}^7 q^{\frac{1}{6}} + \zeta_{54}^{14} q^{\frac{1}{3}} + \zeta_{54} q^{\frac{4}{6}} + \dots \right).$$

$$\text{So } \text{val}_3(f|_{1/3}) = -3, \text{val}_3(f|_{1/9}) = -1.$$

## Examples (contd.)

$$N = 5^2, k = 4, p = 5$$

$$f = q + 4q^2 - 2q^3 + 8q^4 + \dots$$

$$f|_4\left(\frac{1}{5} \ 0\right) = \frac{1}{5} \left( (-4\zeta_5^3 - 3\zeta_5 - 3) q + (-12\zeta_5^2 - 16\zeta_5 - 12) q^2 + \dots \right).$$

$$\text{val}_5(f|_{1/5}) = -1/2.$$

## Examples (contd.)

$$N = 5^2, k = 4, p = 5$$

$$f = q + 4q^2 - 2q^3 + 8q^4 + \dots$$

$$f|_4 \left( \begin{smallmatrix} 1 & 0 \\ 5 & 1 \end{smallmatrix} \right) = \frac{1}{5} \left( (-4\zeta_5^3 - 3\zeta_5 - 3) q + (-12\zeta_5^2 - 16\zeta_5 - 12) q^2 + \dots \right).$$

$$\text{val}_5(f|_{1/5}) = -1/2.$$

$$N = 7^2, k = 4, p = 7$$

$$f = q - 5q^2 + 17q^4 - 45q^8 + \dots$$

$$f|_4 \left( \begin{smallmatrix} 1 & \\ 7 & 1 \end{smallmatrix} \right) = \frac{1}{7} \left( (-2\zeta_7^5 - 4\zeta_7^4 - 6\zeta_7^3 - 8\zeta_7^2 - 3\zeta_7 - 5) q \right. \\ \left. + (-30\zeta_7^5 + 10\zeta_7^4 - 20\zeta_7^3 - 15\zeta_7^2 - 10\zeta_7 - 5) q^2 + \dots \right)$$

$$\text{val}_7(f|_{1/7}) = -1/6.$$

## Examples (contd.)

$$N = 2^8 \cdot 3, k = 2, p = 2$$

$$f = q + q^3 + 4q^7 + \dots$$

$$f|_2 \left( \begin{smallmatrix} 1 & 1 \\ 2 & 3 \end{smallmatrix} \right) = \frac{1}{192} \left( \zeta_{128} q^{\frac{1}{192}} + \zeta_{128}^3 q^{\frac{3}{192}} + \dots \right)$$

$$f|_2 \left( \begin{smallmatrix} 1 & -1 \\ 4 & -3 \end{smallmatrix} \right) = \frac{1}{48} \left( \zeta_{64}^{15} q^{\frac{1}{48}} + \dots \right)$$

$$f|_2 \left( \begin{smallmatrix} 3 & 1 \\ 8 & 3 \end{smallmatrix} \right) = \frac{1}{12} \left( \zeta_{32}^5 q^{\frac{1}{12}} + \dots \right)$$

$$f|_2 \left( \begin{smallmatrix} 5 & -1 \\ 16 & -3 \end{smallmatrix} \right) = \frac{1}{3} \left( 2\zeta_{16}^7 q^{\frac{2}{3}} + \dots \right)$$

$$\text{val}_7(f|_{1/2}) = -6, \text{val}_7(f|_{1/4}) = -4, \text{val}_7(f|_{3/8}) = -2, \text{val}_7(f|_{5/16}) = 1.$$

# The main theorem

## Theorem 1

For a newform  $f$  of weight  $k$  for  $\Gamma_0(N)$ , a prime  $p$ , and a cusp  $c$  of denominator  $L$ , the quantity  $\text{val}_p(f|_c)$  depends **only** on  $f$  and  $\text{val}_p(L)$ .

For  $0 \leq \text{val}_p(L) \leq \frac{\text{val}_p(N)}{2}$ , we have the bounds  $\text{val}_p(f|_c) \geq$

$$-\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L)) + \begin{cases} 0 & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1, \text{val}_p(N) > 2, \\ -\frac{1}{2} & \text{if } \text{val}_p(L) = \frac{1}{2}\text{val}_p(N) = 1, \\ 1 - \frac{1}{2}\text{val}_p(L) & \text{otherwise.} \end{cases}$$

For  $p = 2$ , we get even stronger bounds...

# The main theorem

## Theorem 1 (contd...)

If  $p = 2$  we have the additional stronger bounds.

$$\text{val}_2(f|_c) \geq -\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L))$$

$$+ \begin{cases} 0 & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) = 1, \\ \frac{k}{2} & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) \in \{2, 3, 4\}, \\ \frac{k}{2} + 1 - \frac{1}{4}\text{val}_2(N) & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) > 4, \\ 0 & \text{if } \text{val}_2(L) = 3, \text{val}_2(N) > 6. \end{cases}$$

# The main theorem

## Theorem 1 (contd...)

If  $p = 2$  we have the additional stronger bounds.

$$\text{val}_2(f|_c) \geq -\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L))$$

$$+ \begin{cases} 0 & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) = 1, \\ \frac{k}{2} & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) \in \{2, 3, 4\}, \\ \frac{k}{2} + 1 - \frac{1}{4}\text{val}_2(N) & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) > 4, \\ 0 & \text{if } \text{val}_2(L) = 3, \text{val}_2(N) > 6. \end{cases}$$

- We have checked experimentally that our bounds are **sharp** for newforms associated to elliptic curves and  $p \leq 17$ .

# An application to the Manin constant

## The modularity theorem (Wiles-Taylor, B-C-D-T)

Given an elliptic curve  $E/\mathbb{Q}$  of conductor  $N$ ,

- ( *$E$  is modular*) There exists a newform  $f$  of weight 2 for  $\Gamma_0(N)$  and with integral Fourier coefficients such that  $a_f(p) = p + 1 - |E(\mathbb{F}_p)|$ .
- ( *$E$  has a modular parametrization*) There is a surjection  $\phi: X_0(N)_{\mathbb{Q}} \twoheadrightarrow E$ .

Note:  $\phi$  is not unique, so it is common to normalize  $\phi$  to be **optimal**, that is,  $\deg(\phi)$  to be the least possible.

# An application to the Manin constant

## The modularity theorem (Wiles-Taylor, B-C-D-T)

Given an elliptic curve  $E/\mathbb{Q}$  of conductor  $N$ ,

- ( *$E$  is modular*) There exists a newform  $f$  of weight 2 for  $\Gamma_0(N)$  and with integral Fourier coefficients such that  $a_f(p) = p + 1 - |E(\mathbb{F}_p)|$ .
- ( *$E$  has a modular parametrization*) There is a surjection  $\phi: X_0(N)_{\mathbb{Q}} \twoheadrightarrow E$ .

Note:  $\phi$  is not unique, so it is common to normalize  $\phi$  to be **optimal**, that is,  $\deg(\phi)$  to be the least possible.

- The **Manin constant**  $c_\phi$  is defined by  $\phi^*(\omega_E) = c_\phi \cdot \omega_f$  where  $\omega_E$  is the Néron differential and  $\omega_f = 2\pi i f(z) dz$ .

# An application to the Manin constant

## The modularity theorem (Wiles-Taylor, B-C-D-T)

Given an elliptic curve  $E/\mathbb{Q}$  of conductor  $N$ ,

- ( *$E$  is modular*) There exists a newform  $f$  of weight 2 for  $\Gamma_0(N)$  and with integral Fourier coefficients such that  $a_f(p) = p + 1 - |E(\mathbb{F}_p)|$ .
- ( *$E$  has a modular parametrization*) There is a surjection  $\phi: X_0(N)_{\mathbb{Q}} \twoheadrightarrow E$ .

Note:  $\phi$  is not unique, so it is common to normalize  $\phi$  to be **optimal**, that is,  $\deg(\phi)$  to be the least possible.

- The **Manin constant**  $c_\phi$  is defined by  $\phi^*(\omega_E) = c_\phi \cdot \omega_f$  where  $\omega_E$  is the Néron differential and  $\omega_f = 2\pi i f(z) dz$ .

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

- (Gabber in PhD studies; Edixhoven, 1991)  $c_\phi$  is an integer.
- (Abbes–Ullmo, 1996): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p|N$ .
- Mazur, Raynaud, Agashe–Ribet–Stein,....: Further improvements
- (Cremona): Computationally verified conjecture for all  $N \leq 390000$ .

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

- (Gabber in PhD studies; Edixhoven, 1991)  $c_\phi$  is an integer.
- (Abbes–Ullmo, 1996): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p|N$ .
- Mazur, Raynaud, Agashe–Ribet–Stein,....: Further improvements
- (Cremona): Computationally verified conjecture for all  $N \leq 390000$ .
- **(Česnavičius, 2018): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p^2|N$ . (This implies Manin's conjecture if  $N$  is squarefree)**

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

- (Gabber in PhD studies; Edixhoven, 1991)  $c_\phi$  is an integer.
- (Abbes–Ullmo, 1996): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p|N$ .
- Mazur, Raynaud, Agashe–Ribet–Stein,....: Further improvements
- (Cremona): Computationally verified conjecture for all  $N \leq 390000$ .
- **(Česnavičius, 2018): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p^2|N$ . (This implies Manin's conjecture if  $N$  is squarefree)**

**Recall:**  $v_2(N) \leq 8$ ,  $v_3(N) \leq 5$ ,  $v_p(N) \leq 2$  for  $p > 3$ .

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

- (Gabber in PhD studies; Edixhoven, 1991)  $c_\phi$  is an integer.
- (Abbes–Ullmo, 1996): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p|N$ .
- Mazur, Raynaud, Agashe–Ribet–Stein,....: Further improvements
- (Cremona): Computationally verified conjecture for all  $N \leq 390000$ .
- **(Česnavičius, 2018): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p^2|N$ . (This implies Manin's conjecture if  $N$  is squarefree)**

**Recall:**  $v_2(N) \leq 8$ ,  $v_3(N) \leq 5$ ,  $v_p(N) \leq 2$  for  $p > 3$ .

## Theorem 2

For  $\Gamma_1(N) \subset \Gamma \subset \Gamma_0(N)$ , every surjection  $\phi: (X_\Gamma)_\mathbb{Q} \twoheadrightarrow E$  satisfies  $c_\phi \mid 6 \cdot \deg(\phi)$ , and if  $N$  is cube-free or  $\Gamma = \Gamma_1(N)$ , then even  $c_\phi \mid \deg(\phi)$ .

## Conjecture (Manin, 1972)

If  $\phi$  is optimal then  $c_\phi = \pm 1$ .

- (Gabber in PhD studies; Edixhoven, 1991)  $c_\phi$  is an integer.
- (Abbes–Ullmo, 1996): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p|N$ .
- Mazur, Raynaud, Agashe–Ribet–Stein,....: Further improvements
- (Cremona): Computationally verified conjecture for all  $N \leq 390000$ .
- **(Česnavičius, 2018): If  $\phi$  is optimal and  $p|c_\phi$ , then  $p^2|N$ . (This implies Manin's conjecture if  $N$  is squarefree)**

**Recall:**  $v_2(N) \leq 8$ ,  $v_3(N) \leq 5$ ,  $v_p(N) \leq 2$  for  $p > 3$ .

## Theorem 2

For  $\Gamma_1(N) \subset \Gamma \subset \Gamma_0(N)$ , every surjection  $\phi: (X_\Gamma)_\mathbb{Q} \twoheadrightarrow E$  satisfies  $c_\phi \mid 6 \cdot \deg(\phi)$ , and if  $N$  is cube-free or  $\Gamma = \Gamma_1(N)$ , then even  $c_\phi \mid \deg(\phi)$ .

This is interesting because  $\deg(\phi)$  has little in common with  $N$ . No apparent connection between the conditions  $p^2|N$  and  $p|\deg(\phi)$ .

A very brief sketch of proof of Theorem 2:

- 1 Using Theorem 1, we show that

$$\omega_f \text{ lies in the } \mathbb{Z}\text{-lattice } H^0(X_0(N)_{\mathbb{Z}}, \Omega) \subset H^0(X_0(N)_{\mathbb{Q}}, \Omega^1), \quad (1)$$

where  $\Omega$  denotes the relative dualizing sheaf. (Arithmetic geometric considerations reduce this to certain bounds on the  $p$ -adic valuations of the denominators of the Fourier coefficients of  $f$  at *all* the cusps of  $X_0(N)_{\mathbb{C}}$ . Theorem 1 gives much stronger bounds than needed.)

- 2 Using above, we show that  $\omega_f$  lies in an even *a priori* smaller lattice  $H^0(\mathcal{J}_0(N), \Omega^1)$  that seems otherwise inaccessible. Here  $\mathcal{J}_0(N)$  is the Néron model of the Jacobian  $J_0(N)$ .
- 3 Now Theorem 2 follows from the fact that the composition  $\pi \circ \pi^{\vee} : E \rightarrow J_0(N) \rightarrow E$  is multiplication by  $\deg(\phi)$ .

For the rest of this talk I will focus on the proof of Theorem 1.

Recall Theorem 1:

### Theorem 1

For a newform  $f$  of weight  $k$  for  $\Gamma_0(N)$ , a prime  $p$ , and a cusp  $c$  of denominator  $L$ , the quantity  $\text{val}_p(f|_c)$  depends only on  $f$  and  $\text{val}_p(L)$ .

For  $0 \leq \text{val}_p(L) \leq \frac{\text{val}_p(N)}{2}$ , we have the bounds  $\text{val}_p(f|_c) \geq$

$$-\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L)) + \begin{cases} 0 & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1, \text{val}_p(N) > 2, \\ -\frac{1}{2} & \text{if } \text{val}_p(L) = \frac{1}{2}\text{val}_p(N) = 1, \\ 1 - \frac{1}{2}\text{val}_p(L) & \text{otherwise.} \end{cases}$$

with sharper bounds for  $p = 2$ .

# Fourier expansions and Whittaker models

In order to prove Theorem 1, for a cusp  $\mathfrak{c} = \gamma\infty$  and a prime  $p$ , we want to prove lower bounds on

$$\mathrm{val}_p(f|_{\mathfrak{c}}) := \inf_{n \geq 0} (\mathrm{val}_p(a_f(n; \mathfrak{c})))$$

where

$$(f|_k \gamma)(z) = \sum_{n \geq 0} a_f(n; \mathfrak{c}) q^{\frac{n}{w(\mathfrak{c})}}.$$

# Fourier expansions and Whittaker models

In order to prove Theorem 1, for a cusp  $\mathfrak{c} = \gamma\infty$  and a prime  $p$ , we want to prove lower bounds on

$$\mathrm{val}_p(f|_{\mathfrak{c}}) := \inf_{n \geq 0} (\mathrm{val}_p(a_f(n; \mathfrak{c})))$$

where

$$(f|_k \gamma)(z) = \sum_{n \geq 0} a_f(n; \mathfrak{c}) q^{\frac{n}{w(\mathfrak{c})}}.$$

Fourier coefficients at general cusps are subtle: e.g., the coefficients  $a_f(n; \mathfrak{c})$  are not multiplicative. One way to understand  $a_f(n; \mathfrak{c})$  is via the Whittaker model.

- Let  $\phi_f : \mathrm{GL}_2(\mathbb{A}) \rightarrow \mathbb{C}$  be the automorphic form associated to  $f$  via adelization.

- Let  $\phi_f : \mathrm{GL}_2(\mathbb{A}) \rightarrow \mathbb{C}$  be the automorphic form associated to  $f$  via adelicization.
- $\phi_f$  generates a cuspidal automorphic representation  $\pi = \otimes_v \pi_v$ .

- Let  $\phi_f : \mathrm{GL}_2(\mathbb{A}) \rightarrow \mathbb{C}$  be the automorphic form associated to  $f$  via adelicization.
- $\phi_f$  generates a cuspidal automorphic representation  $\pi = \otimes_v \pi_v$ .
- The **global Whittaker newform**  $W_f(g) = \int_{\mathbb{Q} \backslash \mathbb{A}} \phi_f\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g\right) \psi(-x) dx$  packages together all Fourier coefficients at all cusps. In particular,  $a_f(r; \mathfrak{c}) = W_f(g_{r,\mathfrak{c}})$  for some explicit  $g_{r,\mathfrak{c}} \in \mathrm{GL}_2(\mathbb{A})$ .

- Let  $\phi_f : \mathrm{GL}_2(\mathbb{A}) \rightarrow \mathbb{C}$  be the automorphic form associated to  $f$  via adelization.
- $\phi_f$  generates a cuspidal automorphic representation  $\pi = \otimes_v \pi_v$ .
- The **global Whittaker newform**  $W_f(g) = \int_{\mathbb{Q} \backslash \mathbb{A}} \phi_f\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g\right) \psi(-x) dx$  packages together all Fourier coefficients at all cusps. In particular,  $a_f(r; \mathfrak{c}) = W_f(g_{r, \mathfrak{c}})$  for some explicit  $g_{r, \mathfrak{c}} \in \mathrm{GL}_2(\mathbb{A})$ .
- On the other hand,  $W_f(g) = \prod_v W_{\pi_v}(g_v)$ , where  $W_{\pi_v} : \mathrm{GL}_2(\mathbb{Q}_v) \rightarrow \mathbb{C}$  is the **local Whittaker newform** that depends only on the local representation  $\pi_v$ .

- Let  $\phi_f : \mathrm{GL}_2(\mathbb{A}) \rightarrow \mathbb{C}$  be the automorphic form associated to  $f$  via adelization.
- $\phi_f$  generates a cuspidal automorphic representation  $\pi = \otimes_v \pi_v$ .
- The **global Whittaker newform**  $W_f(g) = \int_{\mathbb{Q} \backslash \mathbb{A}} \phi_f\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} g\right) \psi(-x) dx$  packages together all Fourier coefficients at all cusps. In particular,  $a_f(r; \mathfrak{c}) = W_f(g_{r, \mathfrak{c}})$  for some explicit  $g_{r, \mathfrak{c}} \in \mathrm{GL}_2(\mathbb{A})$ .
- On the other hand,  $W_f(g) = \prod_v W_{\pi_v}(g_v)$ , where  $W_{\pi_v} : \mathrm{GL}_2(\mathbb{Q}_v) \rightarrow \mathbb{C}$  is the **local Whittaker newform** that depends only on the local representation  $\pi_v$ .

## An explicit relation

For a newform  $f$  of weight  $k$  for  $\Gamma_0(N)$ , a prime  $p$ , and a matrix  $\gamma = \begin{pmatrix} a & * \\ L & * \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ ,  $\mathfrak{c} = \frac{a}{L}$ , with  $L^2 | N$ , up to a root of unity:

$$a_f(r; \mathfrak{c}) = a_f(r_0) \left( \frac{r}{r_0 w(\mathfrak{c})} \right)^{k/2} \prod_{q|N} W_{\pi_q} \left( \begin{pmatrix} 0 & \frac{r}{N} \\ 1 & \frac{u_q}{L} \end{pmatrix} \right).$$

where  $r_0$  is the  $N$ -free part of  $r$ , and  $u_q \in \mathbb{Z}_q^\times$ .

- Upshot:** Proving lower bounds for  $\text{val}_p(f|_c)$  reduce to proving lower bounds for  $\text{val}_p \left( W_{\pi_q} \begin{pmatrix} 0 & q^t \\ 1 & \frac{u_q}{q^\ell} \end{pmatrix} \right)$  for primes  $p$  and  $q$  both dividing  $N$ ,  $t \in \mathbb{Z}$ ,  $0 \leq \ell \leq \frac{c(\pi_q)}{2}$ ,  $u_q \in \mathbb{Z}_q^\times$ .

- **Upshot:** Proving lower bounds for  $\text{val}_p(f|_c)$  reduce to proving lower bounds for  $\text{val}_p \left( W_{\pi_q} \begin{pmatrix} 0 & q^t \\ 1 & \frac{u}{q^\ell} \end{pmatrix} \right)$  for primes  $p$  and  $q$  both dividing  $N$ ,  $t \in \mathbb{Z}$ ,  $0 \leq \ell \leq \frac{c(\pi_q)}{2}$ ,  $u \in \mathbb{Z}_q^\times$ .
- Since  $|x|_p = p^{-\text{val}_p(x)}$ , this is a *p-adic analogue* of the local sup-norm question of bounding  $|W_{\pi_q}|_\infty$  in highly ramified cases. (Templier 2014, S. 2016, Assing 2019)
- The values of  $W_{\pi_q}$  at diagonal matrices are well-known, the key point is to access the non-diagonal elements.
- **Remark:** Any matrix  $g$  in  $\text{GL}_2(\mathbb{Q}_q)$  has a double coset representative in  $N(F)gK_0(n)$  of the form  $\begin{pmatrix} 0 & q^t \\ 1 & \frac{u}{q^\ell} \end{pmatrix}$  for  $0 \leq \ell \leq n$ ; local Atkin–Lehner operators halve the range of  $\ell$ .

To prove lower bounds for  $\text{val}_p \left( W_{\pi_q} \begin{pmatrix} 0 & q^t \\ 1 & \frac{u}{q^\ell} \end{pmatrix} \right)$  we refine and extend a method developed for the *sup-norm problem* (S. 2016– 2019, Assing 2018–2019, Assing–Corbett 2019,...).

## The local functional equation (Jacquet–Langlands, 1972)

For a non-archimedean local field  $F$ , an infinite-dimensional representation  $\pi$  of  $\mathrm{GL}_2(F)$ , an element  $W$  in the local Whittaker model of  $\pi$ , and a character  $\mu$  of  $F^\times$ , putting

$$Z(W, s, \mu) = \int_{F^\times} W\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}\right) \mu(y) |y|^{s-\frac{1}{2}} d^\times y$$

$$\frac{Z(W, s, \mu)}{L(s, \pi \otimes \mu)} \varepsilon(s, \pi \otimes \mu) = \frac{Z\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \cdot W, 1-s, \mu^{-1}\right)}{L(1-s, \pi \otimes \mu^{-1})}, \quad (2)$$

Above  $\varepsilon(s, \pi)$  is the local  $\mathrm{GL}_2$   $\varepsilon$ -factor (Jacquet–Langlands).

## The local functional equation (Jacquet–Langlands, 1972)

For a non-archimedean local field  $F$ , an infinite-dimensional representation  $\pi$  of  $\mathrm{GL}_2(F)$ , an element  $W$  in the local Whittaker model of  $\pi$ , and a character  $\mu$  of  $F^\times$ , putting

$$Z(W, s, \mu) = \int_{F^\times} W\left(\begin{pmatrix} y & \\ & 1 \end{pmatrix}\right) \mu(y) |y|^{s-\frac{1}{2}} d^\times y$$

$$\frac{Z(W, s, \mu)}{L(s, \pi \otimes \mu)} \varepsilon(s, \pi \otimes \mu) = \frac{Z\left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \cdot W, 1-s, \mu^{-1}\right)}{L(1-s, \pi \otimes \mu^{-1})}, \quad (2)$$

Above  $\varepsilon(s, \pi)$  is the local  $\mathrm{GL}_2$   $\epsilon$ -factor (Jacquet–Langlands).

Using this, one can formulate a “basic identity” (S, 2016) that writes down  $W_{\pi_q}(g_q)$  as an explicit linear combination of terms involving  $\mathrm{GL}_2$  and  $\mathrm{GL}_1$   $\epsilon$ -factors.

For example, if  $\pi$  is supercuspidal, the basic identity becomes

### The basic identity for supercuspidal reps $\pi$

For a supercuspidal rep  $\pi$  of  $\mathrm{PGL}_2(\mathbb{Q}_q)$ ,  $u \in \mathbb{Z}_q^\times$ , and  $1 \leq \ell \leq \frac{c(\pi)}{2}$ .

$$W_\pi \left( \begin{array}{c} 0 & q^t \\ 1 & \frac{u}{q^\ell} \end{array} \right) = (1 - q^{-1})^{-1} q^{-\frac{\ell}{2}} \sum_{\substack{c(\mu)=\ell \\ c(\mu\pi)=-t}} \varepsilon(1/2, \mu) \varepsilon(1/2, \mu^{-1}\pi) \mu(u). \quad (3)$$

For other representations, the basic identity takes a similar (though slightly more complicated) shape. The resulting formulae were written by me in some cases (S, 2016 - 2018) and in all cases by Assing in his thesis (2019).

For example, if  $\pi$  is supercuspidal, the basic identity becomes

### The basic identity for supercuspidal reps $\pi$

For a supercuspidal rep  $\pi$  of  $\mathrm{PGL}_2(\mathbb{Q}_q)$ ,  $u \in \mathbb{Z}_q^\times$ , and  $1 \leq \ell \leq \frac{c(\pi)}{2}$ .

$$W_\pi \left( \begin{array}{c} 0 & q^t \\ 1 & \frac{u}{q^\ell} \end{array} \right) = (1 - q^{-1})^{-1} q^{-\frac{\ell}{2}} \sum_{\substack{c(\mu)=\ell \\ c(\mu\pi)=-t}} \varepsilon(1/2, \mu) \varepsilon(1/2, \mu^{-1}\pi) \mu(u). \quad (3)$$

For other representations, the basic identity takes a similar (though slightly more complicated) shape. The resulting formulae were written by me in some cases (S, 2016 - 2018) and in all cases by Assing in his thesis (2019).

So we need to solve the problem of computing  **$p$ -adic valuations of  $\varepsilon$ -factors** of representations of  $\mathrm{GL}_r(\mathbb{Q}_q)$  where  $r = 1, 2$ .

## The case $q \neq p$

### Theorem 3

*For a finite extension  $F/\mathbb{Q}_q$ , an infinite-dimensional ramified representation  $\pi$  of  $\mathrm{GL}_2(F)$  associated to a holomorphic newform, and a matrix  $g \in \mathrm{GL}_2(F)$ , we have  $W_\pi(g) \in \overline{\mathbb{Z}} \left[ \frac{1}{q} \right]$ .  
In particular, if  $p \neq q$ , then  $\mathrm{val}_p(W_\pi(g)) \geq 0$ .*

## The case $q \neq p$

### Theorem 3

*For a finite extension  $F/\mathbb{Q}_q$ , an infinite-dimensional ramified representation  $\pi$  of  $\mathrm{GL}_2(F)$  associated to a holomorphic newform, and a matrix  $g \in \mathrm{GL}_2(F)$ , we have  $W_\pi(g) \in \overline{\mathbb{Z}} \left[ \frac{1}{q} \right]$ .  
In particular, if  $p \neq q$ , then  $\mathrm{val}_p(W_\pi(g)) \geq 0$ .*

This relies on a formula for the Whittaker newvector in terms of a family of nonarchimedean  ${}_2F_1$  hypergeometric integrals (Assing 2019; also unpublished works of Templier (2012) and Hu (2016)).

### Sketch of proof of Theorem 3 (assuming above-mentioned formula)

Suppose  $G$  compact group,  $K \subseteq G$  of finite index,  $\mathrm{vol}(K) \in R$ . Let  $f : G \rightarrow R$  be a right- $K$ -invariant function. Then  $\int_G f(g) dg \in R$ .

So we are reduced to the case  $q = p$ .

## The case $q = p$

*So the next problem is:* Let  $F$  be a finite extension of  $\mathbb{Q}_p$ . Understand the  $p$ -adic valuations of  $\varepsilon(1/2, \mu)$  and  $\varepsilon(1/2, \mu \otimes \pi)$  where  $\mu$  is a finite order character of  $F^\times$  and  $\pi$  be an infinite-dimensional, irreducible, unitary representation of  $\mathrm{PGL}_2(F)$ .

## The case $q = p$

*So the next problem is:* Let  $F$  be a finite extension of  $\mathbb{Q}_p$ . Understand the  $p$ -adic valuations of  $\varepsilon(1/2, \mu)$  and  $\varepsilon(1/2, \mu \otimes \pi)$  where  $\mu$  is a finite order character of  $F^\times$  and  $\pi$  be an infinite-dimensional, irreducible, unitary representation of  $\mathrm{PGL}_2(F)$ .

- If  $\pi$  is principal series, we need to also assume that it comes from a global holomorphic newform (otherwise we cannot expect good results).
- Note:  $\varepsilon(1/2, \mu)$  and  $\varepsilon(1/2, \mu \otimes \pi)$  are **algebraic numbers of absolute value 1**, but are **not** necessarily roots of unity.

## The case of $GL_1$

- The  $GL_1$   $\epsilon$ -factors defined by Tate are closely related to classical Gauss sums.
- For a classical Gauss sum, there is a well-known result (Stickelberger's congruence) that gives its  $p$ -adic valuation.

### Theorem 4

For a finite extension  $F/\mathbb{Q}_p$ , and a character  $\chi: F^\times \rightarrow \mathbb{C}^\times$  of finite order,

- ① if  $a(\chi) = 1$ , then,

$$\text{val}_p(\epsilon(\frac{1}{2}, \chi)) = -\frac{[\mathbb{F}_F : \mathbb{F}_p]}{2} + \frac{s(\chi)}{p-1}, \quad 0 \leq s(\chi) \leq (p-1)[\mathbb{F}_F/\mathbb{F}_p];$$

- ② if  $\chi^2 = 1$  or  $a(\chi) > 1$ , then  $\epsilon(\frac{1}{2}, \chi)$  is a root of unity, and so

$$\text{val}_p(\epsilon(\frac{1}{2}, \chi)) = 0.$$

A classification of infinite-dimensional, irreducible, unitary representation of  $GL_2(F)$  and trivial central character.

- ① Principal series representations
- ② Special representations (twists of Steinberg)
- ③ Supercuspidal representations:
  - a Dihedral supercuspidal
  - b **Non-dihedral supercuspidal** (can only occur if  $p = 2$ )

A classification of infinite-dimensional, irreducible, unitary representation of  $GL_2(F)$  and trivial central character.

- ① Principal series representations
- ② Special representations (twists of Steinberg)
- ③ Supercuspidal representations:
  - a Dihedral supercuspidal
  - b **Non-dihedral supercuspidal** (can only occur if  $p = 2$ )

All other cases reduce to  $GL_1$

In cases 1, 2 and 3a, one can write the  $GL_2$   $\varepsilon$ -factor in terms of  $GL_1$   $\varepsilon$ -factors. So the problem here reduces to one we have solved.

A classification of infinite-dimensional, irreducible, unitary representation of  $GL_2(F)$  and trivial central character.

- ① Principal series representations
- ② Special representations (twists of Steinberg)
- ③ Supercuspidal representations:
  - a Dihedral supercuspidal
  - b **Non-dihedral supercuspidal** (can only occur if  $p = 2$ )

All other cases reduce to  $GL_1$

In cases 1, 2 and 3a, one can write the  $GL_2$   $\varepsilon$ -factor in terms of  $GL_1$   $\varepsilon$ -factors. So the problem here reduces to one we have solved.

Analysis of non-dihedral representations

There are exactly 16 representations of Type 3b. Using the Local Langlands correspondence and the basic identity we write down  $W_\pi(g)$  exactly in each case, from which the required bounds follow.

- We now know how to estimate the  $p$ -adic valuations of  $\mathrm{GL}_r$ - $\epsilon$ -factors for  $r = 1, 2$ .
- The basic identity expresses  $W_{\pi_p} \left( \begin{smallmatrix} 0 & p^t \\ 1 & \frac{u_p}{p^\ell} \end{smallmatrix} \right)$  as an explicit finite sum involving the above.
- This allows us to prove our main local result, which gives sharp  $p$ -adic bounds for  $\mathrm{val}_p(W_{\pi_p}(\cdot))$  in all cases.

- We now know how to estimate the  $p$ -adic valuations of  $\mathrm{GL}_r$ - $\epsilon$ -factors for  $r = 1, 2$ .
- The basic identity expresses  $W_{\pi_p} \left( \begin{array}{c} 0 & p^t \\ 1 & \frac{u_p}{p^\ell} \end{array} \right)$  as an explicit finite sum involving the above.
- This allows us to prove our main local result, which gives sharp  $p$ -adic bounds for  $\mathrm{val}_p(W_{\pi_p}(\cdot))$  in all cases.

Here is a special case of our main local result:

## A special case of our local theorem

### Theorem 5

Let  $p$  be odd and  $F/\mathbb{Q}_p$  a finite extension. Let  $\pi$  be a supercuspidal representation of  $\mathrm{GL}_2(F)$ , with trivial central character and  $c(\pi) = n > 2$ . For  $0 \leq \ell \leq n/2$  and  $u \in \mathcal{O}_F$ ,

$$\mathrm{val}_p(W_\pi\left(\begin{pmatrix} 0 & p^\ell \\ 1 & up^{-\ell} \end{pmatrix}\right)) \geq \begin{cases} 0 & \text{if } \ell = 0, 1, \\ [\mathbb{F}_F : \mathbb{F}_p] \left(1 - \frac{\ell}{2}\right) & \text{otherwise.} \end{cases}$$

- Our main local theorem gives such bounds (with *lots* of subcases) covering all representations and conductors.
- If  $p = 2$ , we only do the case  $F = \mathbb{Q}_2$ .
- We get stronger bounds for  $\mathbb{Q}_2$  by exploiting additional parity cancellation in sums of  $\epsilon$ -factors.

Now, Theorem 1 follows as described earlier...

That is, we combine the local bounds on  $\text{val}_p(W_\pi\left(\begin{pmatrix} 0 & p^t \\ 1 & up^{-\ell} \end{pmatrix}\right))$  given by (the general version of) Theorem 5 with

$$a_f(r; \mathfrak{c}) = a_f(r_0) \left(\frac{r}{r_0 w(\mathfrak{c})}\right)^{k/2} \prod_{q|N} W_{\pi_q} \left(\begin{pmatrix} 0 & \frac{r}{N} \\ 1 & \frac{uq}{L} \end{pmatrix}\right)$$

to obtain the sharp lower bounds for  $\text{val}_p(f|_{\mathfrak{c}})$  for holomorphic newforms  $f$  at each cusp  $\mathfrak{c}$ , which is the content of Theorem 1.

## Theorem 1

For a newform  $f$  of weight  $k$  for  $\Gamma_0(N)$ , a prime  $p$ , and a cusp  $\mathfrak{c}$  of denominator  $L$ , the quantity  $\text{val}_p(f|_{\mathfrak{c}})$  depends **only** on  $f$  and  $\text{val}_p(L)$ .

For  $0 \leq \text{val}_p(L) \leq \frac{\text{val}_p(N)}{2}$ , we have the bounds  $\text{val}_p(f|_{\mathfrak{c}}) \geq$

$$-\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L)) + \begin{cases} 0 & \text{if } \text{val}_p(L) = 0, \\ 0 & \text{if } \text{val}_p(L) = 1, \text{val}_p(N) > 2, \\ -\frac{1}{2} & \text{if } \text{val}_p(L) = \frac{1}{2}\text{val}_p(N) = 1, \\ 1 - \frac{1}{2}\text{val}_p(L) & \text{otherwise.} \end{cases}$$

For  $p = 2$ , we get even stronger bounds...

# The main theorem

## Theorem 1 (contd...)

If  $p = 2$  we have the additional stronger bounds.

$$\text{val}_2(f|_c) \geq -\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L))$$

$$+ \begin{cases} 0 & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) = 1, \\ \frac{k}{2} & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) \in \{2, 3, 4\}, \\ \frac{k}{2} + 1 - \frac{1}{4}\text{val}_2(N) & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) > 4, \\ 0 & \text{if } \text{val}_2(L) = 3, \text{val}_2(N) > 6. \end{cases}$$

# The main theorem

## Theorem 1 (contd...)

If  $p = 2$  we have the additional stronger bounds.

$$\text{val}_2(f|_c) \geq -\frac{k}{2} (\text{val}_p(N) - 2\text{val}_p(L))$$

$$+ \begin{cases} 0 & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) = 1, \\ \frac{k}{2} & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) \in \{2, 3, 4\}, \\ \frac{k}{2} + 1 - \frac{1}{4}\text{val}_2(N) & \text{if } \text{val}_2(L) = \frac{1}{2}\text{val}_2(N) > 4, \\ 0 & \text{if } \text{val}_2(L) = 3, \text{val}_2(N) > 6. \end{cases}$$

Thank you!