



ELSEVIER

Contents lists available at ScienceDirect

Journal of Combinatorial Theory, Series B

www.elsevier.com/locate/jctb



Extremal problems for pairs of triangles

Zoltán Füredi^a, Dhruv Mubayi^b, Jason O'Neill^c,
Jacques Verstraëte^c

^a *Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences, P.O. Box 127, Budapest, H-1364, Hungary*

^b *Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, United States of America*

^c *Department of Mathematics, University of California, San Diego, United States of America*



ARTICLE INFO

Article history:

Received 21 December 2020

Available online 24 February 2022

Keywords:

Ordered triple systems

Extremal hypergraphs

Intersecting planar triangle systems

ABSTRACT

A *convex geometric hypergraph* or *cgh* consists of a family of subsets of a strictly convex set of points in the plane. There are eight pairwise nonisomorphic cgh's consisting of two distinct triangles. These were studied at length by Braß [6] (2004) and by Aronov, Dujmović, Morin, Ooms, and da Silveira [2] (2019). We determine the extremal functions exactly for seven of the eight configurations.

The above results are about cyclically ordered hypergraphs. We extend some of them for triangle systems with vertices from a non-convex set. We also solve problems posed by P. Frankl, Holmsen and Kupavskii [15] (2020), in particular, we determine the exact maximum size of an intersecting family of triangles whose vertices come from a set of n points in the plane.

© 2022 Elsevier Inc. All rights reserved.

E-mail addresses: z-furedi@illinois.edu (Z. Füredi), mubayi@uic.edu (D. Mubayi), jmoneill@ucsd.edu (J. O'Neill), jacques@ucsd.edu (J. Verstraëte).

1. Introduction

A *triangle system* is a pair (P, \mathcal{T}) where P is a set of points in the plane in *general position*, i.e., no three collinear, and \mathcal{T} is a set of triangles with vertices from P . (A triangle is a closed set, the convex hull of three points not on a line.) A *convex triangle system* is a triangle system (P, \mathcal{T}) where the elements of P are in strictly convex position. It is convenient to treat P in this case as the vertex set Ω_n of a regular n -gon in the plane, and to consider \mathcal{T} to be a *convex geometric hypergraph* or *cgh* – the vertex set is Ω_n with the clockwise cyclic ordering, and \mathcal{T} is a set of triples from Ω_n called *triangles*. In this language, a cgh \mathcal{S} is *contained* in a cgh \mathcal{T} if there is an injection from the vertex set of \mathcal{S} to the vertex set of \mathcal{T} preserving the cyclic ordering of the vertices and preserving triangles, and we say that a cgh H is *F-free* if H does not contain F . In this paper, we concentrate on extremal problems for pairs of triangles in triangle systems and convex geometric hypergraphs. For the rich history of ordered and convex geometric graph problems and their applications, see [10,18,21,23,25,26,32] and the surveys of Pach [30,31] and Tardos [37], and for convex triangle systems and generalizations, see [7,17,34] and the survey of Braß [6]. On the other hand, the field of extremal hypergraph problems in the convex or geometric setting has fewer results, and statements of general principles in the area are lacking. A natural first step in building such a theory is to solve interesting special cases, and this is one of the goals of this paper.

1.1. Intersecting triangle systems

An old theorem of Hopf and Pannwitz [21] and Sutherland [36] states that the maximum number of line segments between n points in the plane with no two line segments disjoint is n . It is natural to ask for the maximum number of triangles between n points in the plane with no two triangles disjoint. To this end, a triangle system (P, \mathcal{T}) is *intersecting* if any two triangles in \mathcal{T} share at least one point, and *strongly intersecting* if any two triangles in \mathcal{T} share a point in their interior. Intersecting triangle systems are motivated by the Erdős-Ko-Rado Theorem [12], and motivation for considering strong intersection is the well-known theorem of Boros and the first author [5] concerning the *depth* of points. They proved that for every set of n points in the plane, the complete triangle system contains $\frac{2}{9}\binom{n}{3}$ triangles with a common point in their interior (see also Bukh [8], Bukh, Matoušek and Nivasch [9], and Bárány [3], Gromov [20] and Karasev [22] for the d -dimensional analogue). In particular, a strongly intersecting subfamily of size at least $\frac{2}{9}\binom{n}{3}$ exists. P. Frankl, Holmsen and Kupavskii [15] recently determined that the maximum number of triangles in an n -point strongly intersecting convex triangle system is

$$\Delta(n) = \begin{cases} \frac{n(n-1)(n+1)}{24} & \text{if } n \text{ is odd} \\ \frac{n(n-2)(n+2)}{24} & \text{if } n \text{ is even.} \end{cases}$$

In particular, $\Delta(n)/\binom{n}{3} \rightarrow 1/4$ as $n \rightarrow \infty$. The quantity $\Delta(n)$ is also equal to the maximum *depth* of a point in sets of n points in the plane, which can be proved using the *upper bound theorem* for convex polytopes – see Wagner and Welzl [39]. An n -point strongly intersecting convex triangle system of size $\Delta(n)$ is obtained by taking all triangles containing the centroid of Ω_n when n is odd, together with all triangles on one side of each diameter of Ω_n when n is even (these constructions have size $\Delta(n)$, see [4] for instance). For convenience, we let $\mathcal{H}^*(n)$ denote the family of all such convex triangle systems with n points. P. Frankl, Holmsen and Kupavskii posed the following problem (see Problem 1 in [15]):

Problem 1.1. *What is the maximum size, over all point sets of size n , of the largest strongly intersecting triangle system? Is the maximum always at most $(\frac{1}{4} + o(1)) \binom{n}{3}$ as $n \rightarrow \infty$?*

Our first result solves this problem completely for point sets in general position, as follows:

Theorem 1. *Any n -point strongly intersecting triangle system has size at most $\Delta(n)$.*

The short proof of Theorem 1 is given in Section 3. Note that Theorem 1 sharpens and extends the main result of [15] cited above, as $\Delta(n)$ is exactly the size of every convex triangle system in $\mathcal{H}^*(n)$. P. Frankl, Holmsen and Kupavskii further posed the problem of determining the maximum number of triangles in an n -point intersecting convex triangle system if one allows triangles to intersect on the boundary (see Problem 2 in [15]):

Problem 1.2. *What happens if one relaxes the intersecting condition and allows triangles to intersect on the boundary?*

There are a number of different intersection patterns of pairs of triangles in convex triangle systems, depicted below.




Fig. 1. The eight types of triangle pairs in convex triangle systems.

For all of these configurations, Braß [6] has shown the extremal function for convex triangle systems is either $\Theta(n^2)$ or $\Theta(n^3)$; the latter arises precisely when the two triangles have no common interior point. Aronov, Dujmović, Morin, Ooms and da Silveira [2] extensively studied cghs which avoid combinations of the configurations in Fig. 1, and determined many of the order of magnitudes of the associated extremal numbers. An intersecting convex triangle system is precisely a convex triangle system not containing M_1 , and a strongly intersecting convex triangle system is precisely a convex triangle system containing none of D_1, S_1 and M_1 . If \mathcal{F} is a set of convex triangle systems, then we denote by $\text{ex}_\circ(n, \mathcal{F})$ the maximum number of triangles in a convex triangle system not containing any member of \mathcal{F} . In this language, P. Frankl, Holmsen and Kupavskii [15] proved $\text{ex}_\circ(n, \{D_1, S_1, M_1\}) = \Delta(n)$. Problem 1.2 asks for $\text{ex}_\circ(n, \mathcal{F})$ where $\mathcal{F} \subseteq \{D_1, S_1, M_1\}$ and we completely solve this problem using the following theorem:

Theorem 2. For all $n \geq 3$,

$$\text{ex}_\circ(n, F) = \begin{cases} \Delta(n) & \text{if } F = D_1 \\ \Delta(n) + \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor & \text{if } F = S_1 \\ \Delta(n) + \frac{n(n-3)}{2} & \text{if } F = M_1. \end{cases}$$

Furthermore, the extremal constructions for this theorem are classified – see the constructions in Section 2. Using these extremal constructions and Theorem 2, we obtain the exact value of $\text{ex}_\circ(n, \mathcal{F})$ for each $\mathcal{F} \subseteq \{D_1, S_1, M_1\}$:

$$\text{ex}_\circ(n, \mathcal{F}) = \min_{F \in \mathcal{F}} \text{ex}_\circ(n, F).$$

The extremal constructions above are characterized in our proofs in all cases except $\mathcal{F} = \{D_1, M_1\}$.

We also answer Problem 1.2 in the more general context of triangle systems. In this setting, D_1 denotes two triangles on opposite sides of a line and sharing a side – tangent triangles – and S_1 denotes two triangles intersecting in exactly one vertex – touching triangles – whereas M_1 denotes two triangles sharing no points – separated triangles. Theorem 1 and the first two parts of Theorem 2 are an immediate consequence of the following stronger theorem:

Theorem 3. Let $F \in \{D_1, S_1, M_1\}$, and let \mathcal{T} be an n -point triangle system of maximum size not containing F . Then

$$|\mathcal{T}| = \begin{cases} \Delta(n) & \text{if } F = D_1 \\ \Delta(n) + \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor & \text{if } F = S_1 \\ \Delta(n) + \Theta(n^2) & \text{if } F = M_1. \end{cases}$$

The additive term of order n^2 for the case of M_1 in Theorem 3 arises from a geometric theorem of Valtr on *avoiding line segments* in the plane. We believe that the value of $\text{ex}_\cup(n, M_1)$ should determine the maximum for n -point intersecting triangle systems:

Conjecture 1. *For all $n \geq 3$, if \mathcal{T} is an n -point intersecting triangle system, then $|\mathcal{T}| \leq \text{ex}_\cup(n, M_1)$.*

For the above configurations $F \in \{D_1, S_1, M_1\}$, the extremal functions $\text{ex}_\cup(n, F)$ and the maximum number of triangles in an F -free n -point triangle system are equal (almost equal). This is quite exceptional. E.g., one can find a self-intersecting length three path, P_3 , in a convex geometric graph with $\Omega(n)$ edges, while for the general not necessarily convex case Pach, Pinchasi, Tardos, and Tóth [33] showed that there exists P_3 -free geometric graphs with $\Theta(n \log n)$ edges.

1.2. The five configurations in the $\Theta(n^2)$ range

Braß [6] has shown that the five configurations whose extremal function is in the $\Theta(n^2)$ range are S_2, S_3, M_2, M_3 and D_2 . In this section, we present our results for $\text{ex}_\cup(n, F)$ when $F \in \{S_3, M_2, M_3\}$ and our bounds for $F \in \{S_2, D_2\}$. The extremal function for M_3 was determined exactly in [16] and we repeat its proof from [16] for convenience. We also determine the exact extremal function for M_2 and S_3 when n is even:

Theorem 4.

$$\text{ex}_\cup(n, F) = \begin{cases} \binom{n}{3} - \binom{n-3}{3} & \text{if } F = M_3 \text{ and } n \geq 3. \\ \binom{n}{2} - 2 & \text{if } F = M_2 \text{ and } n \geq 7. \\ \frac{n(n-2)}{2} & \text{if } F = S_3 \text{ and } n \geq 4 \text{ is even.} \end{cases}$$

For S_3 when n is odd, there are several constructions which obtain the lower bound $\text{ex}_\cup(n, S_3) \geq \frac{(n-1)(n-2)}{2} + 1$ (see Construction 6 in Section 2), but we have not proved that this bound is sharp. We leave the following open problem:

Problem 1.3. *Prove $\text{ex}_\cup(n, S_3) = (n-1)(n-2)/2 + 1$ when $n \geq 5$ is odd, and characterize the extremal S_3 -free convex geometric hypergraphs.*

The configurations S_2 and D_2 appear to be the most difficult to handle.

Theorem 5. *For $n \geq 3$,*

$$\left\lfloor \frac{n^2}{4} \right\rfloor - 1 \leq \text{ex}_\cup(n, S_2) \leq \frac{23}{64}n^2.$$

We believe that the lower bound in this theorem is tight.

Conjecture 2. For all $n \geq 5$, $\text{ex}_\circ(n, S_2) = \lfloor n^2/4 \rfloor - 1$.

Theorem 6. For $n \geq 3$,









$$\frac{4}{9} \binom{n}{2} - O(n) \leq \text{ex}_\circ(n, D_2) \leq \frac{2n^2 - 3n}{9}.$$

The lower bound is due to Damásdi and N. Frankl [11] who solved our conjecture from an earlier draft of this paper and determined $\lim_{n \rightarrow \infty} \text{ex}_\circ(n, D_2)/\binom{n}{2}$. Even more, they showed that equality holds for all $n \equiv 6 \pmod 9$ and gave an independent proof for our upper bound. Beside the upper bound we present a lower bound $\frac{3}{7} \binom{n}{2} - O(n) \leq \text{ex}_\circ(n, D_2)$ in Construction 8 using a quite different method.

1.3. Summary of results

We summarize the results for $\text{ex}_\circ(n, F)$ in this paper in Table 1. For S_2 and D_2 , we only have bounds on the extremal function, and write $[a, b]$ in the table to denote $a \leq \text{ex}_\circ(n, F) \leq b$. We conjecture $\text{ex}_\circ(n, S_2) = \lfloor n^2/4 \rfloor - 1$. The constructions refer to those numbered 1 – 8 in Section 2.

Table 1
Summarized results for $\text{ex}_\circ(n, F)$.

F	$\text{ex}_\circ(n, F)$	Construction	F	Bounds on $\text{ex}_\circ(n, F)$	Construction
 M_1	$\Delta(n) + \frac{n(n-3)}{2}$	3	 S_1	$\Delta(n) + \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor$	2
 M_2	$\binom{n}{2} - 2$	5	 S_2	$[\lfloor \frac{n^2}{4} \rfloor - 1, \frac{23n^2}{64}]$	7
 M_3	$\binom{n}{3} - \binom{n-3}{3}$	4	 D_1	$\Delta(n)$	1
 S_3	$\frac{n(n-2)}{2}$ for n even	6	 D_2	$\frac{2n^2 - 3n}{9}$ for $n \equiv 6 \pmod 9$	8

1.4. Organization

Constructions of F -free convex triangle systems which give lower bounds for the theorems in this paper are in Section 2, Constructions 1 – 8. Sections 3, 4, 5 contain the proofs of our results for D_1 , S_1 and M_1 , respectively (i.e. the proofs of Theorems 2 and 3). Section 6 – 10 contain the proofs of our results concerning the configurations S_2, S_3, M_2, M_3, D_2 . Concluding remarks and further questions are in Section 11.

1.5. Notation

We refer to a set of triples from a set Ω_n of n vertices of a regular n -gon as a *convex triangle system*. It is convenient also to refer to this as a *convex geometric hypergraph* or *cgh*, where the triangles are considered as triples in $\binom{\Omega_n}{3}$, and the vertices of Ω_n are cyclically ordered in the clockwise direction, say $v_0 < v_1 < \dots < v_{n-1} < v_0$. In this case, we consider the subscripts modulo n . A cgh F is *contained* in a cgh H if there is an injection from $V(F)$ to $V(H)$ preserving the cyclic ordering of the vertices and preserving triangles, and we say that H is *F -free* if H does not contain F as a subhypergraph. The extremal function $\text{ex}_\odot(n, F)$ denotes the maximum number of triangles in an F -free cgh on Ω_n . Given $H \subset \binom{\Omega_n}{3}$ and $A \subseteq \Omega_n$, let $d_H(A) = |\{e \in H : A \subset e\}|$ be the *degree* of A in H ; we write $d_H(u, v)$ when $A = \{u, v\}$ and $d_H(v_i)$ when $A = \{v_i\}$. Further, when $A = \{v_i\}$, we may form an ordered graph from H by considering $G_i = \{\{u, w\} : \{v_i, u, w\} \in H\}$ – this is the *link graph* of v_i in H with vertex set $\{v_{i+1}, v_{i+2}, \dots, v_{i-1}\}$ with the natural ordering. Let $\partial H = \{\{u, v\} : \exists e \in H, \{u, v\} \subset e\}$ denote the *shadow* of H . For functions $f, g : \mathbb{N} \rightarrow \mathbb{R}^+$, we write $f = o(g)$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 0$, and $f = O(g)$ if there is $c > 0$ such that $f(n) \leq cg(n)$ for all $n \in \mathbb{N}$. If $f = O(g)$ and $g = O(f)$, we write $f = \Theta(g)$.

2. Constructions

Construction 1 (D_1, S_1 and M_1 -free cghs). For $n \geq 3$ odd, let the class of cghs $\mathcal{H}^*(n)$ comprise the single cgh consisting of triangles which contain in their interior the centroid of Ω_n . For $n \geq 4$ even, each $H \in \mathcal{H}^*(n)$ consists of all triangles which contain the centroid of Ω_n and, for each diameter $\{v_i, v_{i+n/2}\}$ of Ω_n , we either add all triangles $\{v_i, v_j, v_{i+n/2}\}$ where $v_i < v_j < v_{i+n/2} < v_i$, or all triangles $\{v_i, v_j, v_{i+n/2}\}$ where $v_{i+n/2} < v_j < v_i < v_{i+n/2}$. It is not hard to show that each element $H \in \mathcal{H}^*(n)$ has size $\triangle(n)$ – see [4]. Each $H \in \mathcal{H}^*(n)$ is strongly intersecting, so

$$\text{ex}_\odot(n, D_1) \geq \text{ex}_\odot(n, \{D_1, S_1, M_1\}) \geq \triangle(n). \quad (1)$$

Construction 2 (S_1 and M_1 -free cghs). For $n \geq 3$ odd, each cgh in $\mathcal{H}^+(n)$ is obtained by adding for some $i < n$ to any cgh in $\mathcal{H}^*(n)$ all triangles containing a pair $\{v_{i+j}, v_{i+j+(n-1)/2}\}$ for $0 \leq j \leq (n-3)/2$ (left diagram in Fig. 2). For $n \geq 4$ even, each

$H \in \mathcal{H}^+(n)$ consists of all triangles containing the centroid of Ω_n in their interior or on their boundary (right diagram in Fig. 2).




Fig. 2. Construction of $\mathcal{H}^+(n)$.

By inspection, each $H \in \mathcal{H}^+(n)$ is S_1 -free and M_1 -free. Moreover, if n is odd, then $|H| = \Delta(n) + (n-1)(n-3)/4$ whereas if n is even, then $|H| = \Delta(n) + n(n-2)/4$. We obtain

$$\text{ex}_{\odot}(n, S_1) \geq \text{ex}_{\odot}(n, \{S_1, M_1\}) \geq \Delta(n) + \lfloor \frac{n}{2} \rfloor \lfloor \frac{n-2}{2} \rfloor. \quad (2)$$

Construction 3 (M_1 -free cghs). For $n \geq 3$ odd, the unique cgh in $\mathcal{H}^{++}(n)$ is obtained by adding all triangles containing a pair $\{v_i, v_{i+(n-1)/2}\}$ to the cgh in $\mathcal{H}^*(n)$ (left diagram in Fig. 3). For $n \geq 4$ even, $\mathcal{H}^{++}(n)$ is obtained by adding all triangles containing a diameter of Ω_n , plus all triangles containing a pair from a set of $n/2$ pairwise intersecting pairs of the form $\{v_i, v_{i+n/2-1}\}$ to any cgh in $\mathcal{H}^*(n)$ (right diagram in Fig. 3). Every cgh in $\mathcal{H}^{++}(n)$ is M_1 -free, and has size $\Delta(n) + n(n-3)/2$.




Fig. 3. Construction of $\mathcal{H}^{++}(n)$.

Construction 4 (M_3 -free cghs). An extremal M_3 -free construction is simply to take all $n(n-3)$ triangles which contain a pair of cyclically consecutive vertices of Ω_n plus the set of all $\binom{n-4}{2}$ triangles without consecutive elements and containing a fixed vertex v_0 . It turns out this is not the only M_3 -free construction with that many triangles: we may remove any triangle $\{v_0, v_{2k+1}, v_{2k+3}\}$ and add $\{v_{2k}, v_{2k+2}, v_{2k+4}\}$ when $2k+4 < n$ to obtain many different M_3 -free extremal constructions.

Construction 5 (M_2 -free cghs). An M_2 -free construction on Ω_n is obtained by taking all triangles containing a fixed vertex, plus all n triangles of three cyclically consecutive vertices.

The restriction $n \geq 7$ is necessary in Theorem 4, since for $n = 6$, the only copies of M_2 on Ω_6 are the triangles $\{v_0, v_1, v_3\}$, $\{v_0, v_2, v_3\}$, $\{v_0, v_1, v_4\}$, $\{v_0, v_3, v_4\}$, $\{v_0, v_2, v_5\}$, $\{v_0, v_3, v_5\}$ with their corresponding complements. As such, removing exactly one member from each copy of M_2 from the complete cgh on Ω_6 gives an M_2 -free cgh H with $14 = \binom{n}{2} - 1$ triangles. It is likely the case that the set of triangles containing v_0 plus the set of triangles of consecutive vertices in Ω_n is the unique extremal M_2 -free example up to isomorphism for $n \geq 8$. For $n = 7$, we may take all seven cyclically consecutive triangles, the triangle $\{v_1, v_3, v_6\}$, and all triangles which contain v_0 besides the triangle $\{v_0, v_4, v_5\}$. Similarly, when $n = 7$, we may also take all seven cyclically consecutive triangles, the triangles $\{v_1, v_3, v_6\}$ and $\{v_1, v_4, v_6\}$, and all triangles which contain v_0 besides the triangles $\{v_0, v_4, v_5\}$ and $\{v_0, v_2, v_3\}$.

Construction 6 (S_3 -free cghs). For even $n \geq 4$, let

$$H_0 := \left\{ \{v_{2i-1}, v_{2i}, v\} \in \binom{\Omega_n}{3} : 1 \leq i \leq n/2, v \in \Omega_n \setminus \{v_{2i-1}, v_{2i}\} \right\}.$$

By inspection, H_0 is S_3 -free and has $n(n-2)/2$ triangles. For n odd, let H_1 have vertex set $\{v_0, v_1, \dots, v_{n-1}\}$ and add to a copy of H_0 on $\{v_1, v_2, \dots, v_{n-1}\}$ all triangles $\{v_0, v_{2i-1}, v_{2i}\}$ where $1 \leq i \leq (n-1)/2$ as well as $\{v_{n-1}, v_0, v_1\}$. Then H_1 is S_3 -free and $|H_1| = (n-1)(n-2)/2 + 1$.

Construction 7 (S_2 -free cghs). A construction demonstrating the lower bound is to split Ω_n into two intervals A and B , and to take all triangles which contain a point from A and a pair of consecutive points in B . We also add all triangles containing three consecutive points in B . This configuration has $|A|(|B|-1) + |B|-2 = (|A|+1)(|B|-1) - 1$ triangles and does not contain S_2 . If $|A| = \lceil n/2 \rceil - 1$ and $|B| = \lfloor n/2 \rfloor + 1$ then this configuration has $\lfloor n^2/4 \rfloor - 1$ triangles.

Construction 8 (D_2 -free cghs). For a lower bound on $\text{ex}_\cup(n, D_2)$, start with an $S(n, 15, 2)$ design – Wilson [40] proved these exist whenever n is large enough and satisfies the requisite divisibility conditions, i.e., $\binom{n}{2}/\binom{15}{2}$ is an integer, and $n \equiv 1 \pmod{14}$, i.e., $n \equiv 1, 15, 85, 141 \pmod{210}$. The construction is as follows: decompose the $E(K_n)$ into $\binom{n}{2}/\binom{15}{2}$ complete K_{15} 's. Each corresponds to a convex 15-gon with vertex set $V = \{w_1, w_2, \dots, w_{15}\}$. Decompose each K_{15} into fifteen triangulations of a convex pentagon $\{w_i, w_{i+1}, w_{i+6}, w_{i+8}, w_{i+11}\}$ with diagonals $\{w_i, w_{i+6}\}$ and $\{w_i, w_{i+8}\}$ (indices are mod 15). The lengths of the sides are 1, 5, 2, 3, 4 and the diagonals are 6 and 7, so this is indeed a decomposition with 45 triangles. This construction has size exactly

$$45 \cdot \frac{\binom{n}{2}}{\binom{15}{2}} = \frac{3}{7} \binom{n}{2}$$

whenever $n \equiv 1, 15, 85, 141 \pmod{210}$, and gives a construction of size $\frac{3}{7}\binom{n}{2} - O(n)$ for all n .

3. Proof of Theorem 3: tangent triangles, D_1

A *directed 3-cycle* in a tournament is a triple $\{x, y, z\}$ with $x \rightarrow y \rightarrow z \rightarrow x$. Let $T(n)$ be the maximum number of directed 3-cycles in an n -vertex *tournament*. It was shown by Moon [29] (see also pages 42–44 in Erdős and Spencer [13]) that $T(n) = \Delta(n)$ for $n \geq 3$. To see this, every tournament with n vertices of outdegrees d_1, \dots, d_n has exactly $\binom{n}{3} - \sum_{i=1}^n \binom{d_i}{2}$ directed 3-cycles. This is maximized (only) when the outdegrees are as equal as possible. If n is odd, then all $d_i = (n-1)/2$ while if n is even, half of the d_i are $(n-2)/2$ and the other half are $n/2$. These tournaments are called *almost regular*. Tournaments with these outdegrees can easily be constructed and moreover there are plenty of them when n is large. A short calculation gives the required

$$T(n) = \Delta(n).$$

A directed 3-cycle $\{x, y, z\}$ in the plane with $x \rightarrow y \rightarrow z \rightarrow x$ is oriented *clockwise* if z is in the half plane to the right when traversing the segment $[xy]$ from x to y . If $\{x, y, z\}$ is not oriented clockwise, then it is oriented *counterclockwise*.

Proof of Theorem 1. Let P be a set of n points in the plane with no three collinear, and let \mathcal{T} be a D_1 -free family of triangles on P , and let H be the corresponding 3-uniform hypergraph with vertex set P . We will prove that $|\mathcal{T}| \leq T(n)$, which gives Theorem 1.

We define an orientation for each edge $\{x, y\} \in \partial H$ as follows. Consider any triangle $\{x, y, z\} \in \mathcal{T}$. If the orientation of the 3-cycle $\{x, y, z\}$ is clockwise then orient the edge $\{x, y\}$ as $x \rightarrow y$, and $y \rightarrow x$ otherwise. The main observation is that the orientation of $\{x, y\}$ is uniquely determined. If $\{x, y\}$ belongs to an $\{x, y, z\}$ 3-cycle oriented clockwise and to an $\{x, y, z'\}$ 3-cycle oriented counterclockwise, then the two corresponding triangles form D_1 . We conclude that $|H|$ is at most the number of directed 3-cycles in an orientation of a subgraph of K_n which is at most $T(n)$ as required. \square

Extremal families. The previous proof shows that $|H| = \Delta(n)$ is only possible if the orientation of the 3-cycles of ∂H is an almost regular tournament. There is a one to one correspondence between extremal D_1 -free cghs (or a D_1 -free triangle system in general) and almost regular tournaments.

3.1. Extremal $\{D_1, S_1\}$ -free cghs

Suppose that a cgh H is D_1 -free and also S_1 -free with $|H| = \Delta(n)$. Then ∂H is an almost regular tournament and H is obtained as the family of oriented three-cycles in ∂H . We claim that more is true, $H \in \mathcal{H}^*(n)$ as described in Construction 1.

First we show that all directed 3-cycles in the tournament have the same orientation. As a first step, we prove that if two triangles in H have some common vertices then their corresponding 3-cycles have the same orientation. This is obviously true when they have a common edge because H is D_1 -free. Consider first the case when two triangles $T_1, T_2 \in \mathcal{T}$ share a vertex v_1 and have opposite orientations. Then they can form an S_1 (which we excluded), or an S_2 , or an S_3 .

If they form S_2 , say $v_1 < v_2 < \dots < v_5 < v_1$ and the two 3-cycles are oriented as $v_1 \rightarrow v_2 \rightarrow v_5 \rightarrow v_1$ and $v_1 \rightarrow v_4 \rightarrow v_3 \rightarrow v_1$, then we proceed as follows. Consider the edge $\{v_4, v_5\}$. Observe that $v_5 \rightarrow v_4$, otherwise the triangles $\{v_1, v_4, v_5\}$ and $\{v_1, v_3, v_4\}$ form a D_1 . A similar argument shows $v_3 \rightarrow v_2$. Consider the edge $\{v_2, v_4\}$. Now $v_2 \rightarrow v_4$, otherwise the triangles $\{v_1, v_2, v_5\}$ and $\{v_2, v_4, v_5\}$ form a D_1 . But then we have found a triangle $\{v_2, v_4, v_3\}$ which forms an S_1 with $\{v_1, v_2, v_5\}$. So T_1 and T_2 cannot form an S_2 .

If T_1 and T_2 form an S_3 , say $v_1 < \dots < v_5 < v_1$ and the two 3-cycles are oriented as $v_1 \rightarrow v_2 \rightarrow v_4 \rightarrow v_1$ and $v_1 \rightarrow v_5 \rightarrow v_3 \rightarrow v_1$ then we proceed the same way. Consider the edge $\{v_4, v_5\}$. Observe that $v_4 \rightarrow v_5$, otherwise the triangles $\{v_1, v_5, v_4\}$ and $\{v_1, v_2, v_4\}$ form a D_1 . A similar argument shows $v_3 \rightarrow v_2$. Consider the edge $\{v_3, v_4\}$. Then $v_3 \rightarrow v_4$, otherwise the triangles $\{v_1, v_2, v_4\}$ and $\{v_2, v_4, v_3\}$ form a D_1 . But then we have found the triangle $\{v_3, v_4, v_5\}$ which forms a D_1 with $\{v_1, v_5, v_3\}$. So T_1 and T_2 cannot form an S_3 .

The above argument implies that the vertex sets $X := \{x \in e \in H, e \text{ is oriented clockwise}\}$ and $Y := \{y \in e \in H, e \text{ is oriented counterclockwise}\}$ are disjoint. So every triangle $e \in H$ is contained entirely in X or in Y . This gives

$$|H| \leq T(|X|) + T(|Y|) < T(n),$$

a contradiction.

From now on, we may suppose that each directed 3-cycle of ∂H is oriented clockwise. This implies that for $v_i < v_j < v_k < v_i$ we have $v_k \rightarrow v_i$ if $v_j \rightarrow v_i$. Indeed, each orientation of a triangle comes from a directed 3-cycle, so in case of $v_i \rightarrow v_k$ and $v_j \rightarrow v_i$ we get two 3-cycles $\{v_i, v_k, v_{k'}\}$ and $\{v_j, v_i, v_{j'}\}$ oriented clockwise so $v_i < v_{j'} < v_j < v_k < v_{k'} < v_i$, and these two triangles form an S_1 , a contradiction. Summarizing, each v_i has out-edges $v_i \rightarrow v_j$ for $i < j \leq i + \lfloor (n-1)/2 \rfloor$ and in-edges $v_j \rightarrow v_i$ for $i - \lfloor (n-1)/2 \rfloor \leq j < i$, in other words $H \in \mathcal{H}^*(n)$. \square

4. Proof of Theorems 2 and 3: touching triangles, S_1

4.1. Proof of Theorem 2 for S_1

We will prove that $\text{ex}_\cup(n, S_1) \leq \text{ex}_\cup(n, D_1) + \lfloor n/2 \rfloor \lfloor (n-2)/2 \rfloor$ which via Theorem 2 for D_1 gives the upper bound in Theorem 2 for S_1 . For any cgh H , define a graph $G := G(H)$ with $G \subset \partial H$, called the D_1 -graph of H as the set of $\{u, v\}$ for which there are $x, y \in \Omega_n$ with $u < x < v < y < u$ and triangles $\{u, x, v\}$ and $\{u, y, v\}$ in H . In other

words, $\{u, v\}$ has triangles on both sides. This definition can be naturally extended to triangle systems (P, \mathcal{T}) .

Let $H \subset \binom{\Omega_n}{3}$ be a cgh containing no copy of S_1 . We claim that the D_1 -graph G is a matching. Otherwise, if there are $\{u, v\}$ and $\{v, w\}$ in G , then there are $x, y \in \Omega_n$ with $u < x < v < y < w < u$ and triangles $\{u, x, v\}$ and $\{v, y, w\}$ in H which form S_1 . We obtain $|G| \leq \lfloor n/2 \rfloor$. For each $\{u, v\} \in G$, delete all triangles containing $\{u, v\}$ on the side that has fewer triangles (if both sides have the same number of triangles then pick a side arbitrarily). Altogether we delete at most $|G| \lfloor (n-2)/2 \rfloor \leq \lfloor n/2 \rfloor \lfloor (n-2)/2 \rfloor$ triangles. Let $H' \subset H$ be the set of triangles that remain. Since H' is D_1 -free, $|H'| \leq \text{ex}_\cup(n, D_1)$, and we are done.

If H is an extremal S_1 -free cgh, then $|G| = \lfloor n/2 \rfloor$ and each edge $\{u, v\} \in G$ is contained in at least $\lfloor (n-2)/2 \rfloor$ triangles $\{u, v, w\}$ with $u < w < v < u$ and another at least $\lfloor (n-2)/2 \rfloor$ triangles $\{u, v, z\}$ with $u < v < z < u$. This is only possible if the segments representing the edges of G are pairwise crossing each other inside Ω_n . In case of even n we have that G consists of the $n/2$ diameters $\{v_i, v_{i+n/2}\}$ of Ω_n , in case of odd n we may suppose that $G = \{\{v_j, v_{j+(n-1)/2}\} : 0 \leq j \leq (n-3)/2\}$. Since H' is an extremal $\{D_1, S_1\}$ -free cgh the results of subsection 3.1 yield that $H' \in \mathcal{H}^*(n)$. The triangles from $H \setminus H'$ can be added to H' only as described in Construction 2, and this yields $H \in \mathcal{H}^+(n)$. \square

4.2. A geometric lemma about D_1 -edges in S_1 -free triangle systems

Recall that a segment $[ab]$ (with $a, b \in P$) is a D_1 -edge in the triangle system (P, \mathcal{T}) if there are triangles from \mathcal{T} on both sides, i.e., $\exists c^-, c^+ \in P$ such that c^- and c^+ are separated by the line $\ell(ab)$ and $\{a, b, c^-\}, \{a, b, c^+\} \in \mathcal{T}$.

Lemma 4.1. *Let \mathcal{T} be a triangle system with point set P . Suppose that $[ab]$, $[bc]$, and $[cd]$ are distinct D_1 -segments in \mathcal{T} (so $a = d$ is not excluded). Then \mathcal{T} contains an S_1 configuration.*

Proof. The lines $\ell(ab)$, $\ell(bc)$, and $\ell(cd)$ cut the plane into seven open regions, unless $\ell(ab) \parallel \ell(cd)$ when we get only six regions. Let T be the triangle these lines enclose (in the case of six regions T is one of the infinite three-sided strips). Let $H(xy)$ denote the open half plane with boundary line xy tangent to T but disjoint from its interior. Since $\{a, b\}$ is a D_1 -edge there exists a triangle $\{a, b, c^-\} \in \mathcal{T}$ where c^- is in the open half plane $H(ab)$, and there exists a triangle $\{b, c, y\} \in \mathcal{T}$ with $y \in H(bc)$. These two triangles form an S_1 configuration unless $c^-, y \in B := (H(ab) \cup \ell(ab)) \cap H(bc)$. Consider a third triangle $\{c, d, b^-\} \in \mathcal{T}$ where $b^- \in H(cd)$. Since this half plane is separated from $\{b, c, y\}$ by $\ell(cd)$ (except both contain c in their boundaries), so $\{c, d, b^-\}$ and $\{b, c, y\}$ form an S_1 . \square

4.3. A removal lemma concerning S_1 -free triangle systems

We prove Theorem 3 for S_1 in the following stronger form.

Theorem 7. *Let $n \geq 3$, and let \mathcal{T} be an n -point triangle system. If \mathcal{T} is S_1 -free then there exists a subfamily $\mathcal{T}' \subset \mathcal{T}$ which is D_1 -free and*

$$|\mathcal{T}| \leq |\mathcal{T}'| + \left\lfloor \frac{n}{2} \right\rfloor \left\lfloor \frac{n-2}{2} \right\rfloor.$$

Since $|\mathcal{T}'| \leq \Delta(n)$ by Theorem 1, one obtains the desired upper bound for $|\mathcal{T}|$.

Recall that the D_1 -graph of \mathcal{T} is a graph G with vertex set P and its edges are the D_1 -segments. For $e \in G$, let $\mathcal{T}(e)$ be the set of triangles from \mathcal{T} containing e , and let $\text{del}(\{e\})$ be the minimum number of triangles $e \cup \{x\} \in \mathcal{T}$ on one side of $\ell(e)$. Obviously, $\text{del}(\{e\}) \leq (1/2)|\mathcal{T}(e)| \leq \lfloor (n-2)/2 \rfloor$. We extend this definition for any set of edges, $\mathcal{T}(F)$ is the set of triangles from \mathcal{T} containing an edge $e \in F$, and $\text{del}(E(F))$ is the minimum number of triangles $e \cup \{x\} \in \mathcal{T}$, $e \in F$ such that removing those triangles from \mathcal{T} we eliminate all D_1 -edges of F . Our aim is to prove that $\text{del}(E(G)) \leq \lfloor n/2 \rfloor \lfloor (n-2)/2 \rfloor$. We also show that for $n \neq 5$ in case of equality G is either a matching of size $\lfloor n/2 \rfloor$, or a matching of size $(n-3)/2$ and a path of length two. We conjecture that the latter case cannot happen for n sufficiently large.

Since \mathcal{T} is S_1 -free, Lemma 4.1 implies that G contains no path of length three and in particular G does not contain a cycle. Thus G is a starforest.

Claim 4.1. *Suppose that $\{e, f\} \subset G$ is a two-edge component of the D_1 -graph G . Then $\text{del}(\{e, f\}) \leq \lfloor (n-2)/2 \rfloor$.*

Proof. Given the two-edge component $\{e, f\} \subset G$, there exists a $w \in P$ with $w = e \cap f$. Let $\delta = 1$ if $e \cup f \in \mathcal{T}$, and $\delta = 0$ otherwise. We assume e, f are as shown in Fig. 4, i.e., the lines $\ell(e)$ and $\ell(f)$ cut the plane into four open regions A, B, C, D (B is disjoint to $e \cup f$, the boundary of D contains both, etc.). If any of the dotted triangles are in \mathcal{T} , then we get a copy of S_1 , as we have seen this in the proof of Lemma 4.1. More formally, we get, e.g., if $e \cup \{x\} \in \mathcal{T}$, then either $e \cup \{x\} = e \cup f$ or $x \in B \cup C$. For $X \in \{A, B, C, D\}$, let e_X be the number of $x \in X$ such that $e \cup \{x\} \in \mathcal{T}$ and f_X be the number of $x \in X$ such that $f \cup \{x\} \in \mathcal{T}$. We have $e_A, e_D = 0$ and $f_C, f_D = 0$. Observe that $e \cup \{x\}, f \cup \{x\} \in \mathcal{T}$ is not possible for $x \in B$, else we get a D_1 -edge $\{w, x\}$, contradicting the fact that $\{e, f\}$ is a component of G . We obtain

$$f_A + e_B + f_B + e_C \leq |P| - |\{e \cup f\}| \leq n - 3. \quad (3)$$

There are four possibilities to delete triangles from \mathcal{T} to make $\{e, f\}$ non- D_1 -edges, namely we can eliminate all triangles $e \cup \{x\}$ with $x \in A \cup B$ or all such triangles from




Fig. 4. A two-edge component of G as discussed in Claim 4.1.

the other side of $\ell(e)$, and there are two sides of $\ell(f)$ as well. We get four inequalities for $\text{del}(\{e, f\})$.

$$\begin{aligned}\text{del}(\{e, f\}) &\leq e_B + f_B \\ \text{del}(\{e, f\}) &\leq e_B + (f_A + \delta) \\ \text{del}(\{e, f\}) &\leq (e_C + \delta) + f_B \\ \text{del}(\{e, f\}) &\leq (e_C + \delta) + f_A.\end{aligned}$$

Summing these and using (3) we get $4 \text{del}(\{e, f\}) \leq 2n - 6 + 3\delta \leq 2n - 3$. This gives $\text{del}(\{e, f\}) \leq \lfloor (2n - 3)/4 \rfloor = \lfloor (n - 2)/2 \rfloor$ and we are done. \square

Claim 4.2. Suppose that $F \subset G$ is a component of the D_1 -graph G , a star with $s \geq 3$ edges. Then $\text{del}(E(F)) \leq \lfloor (n - 1)/2 \rfloor$.

Proof. We will prove the stronger statement $|\mathcal{T}(F)| \leq n - 1$. Suppose that the edges of F are $\{w, v_1\}, \{w, v_2\}, \dots, \{w, v_s\}$. We claim that for any vertex $x \in P \setminus \{w\}$ an (open) half plane with boundary line $\ell(wx)$ can contain only at most one triangle from \mathcal{T} of the form $\{w, x, v_i\}$. Indeed, if there is another such triangle $\{w, x, v_j\}$ and, say, $\angle(xwv_i) < \angle(xwv_j)$ then there is another vertex $z \in P$ such that $\{w, v_j, z\} \in \mathcal{T}$ and it is separated from $\{w, x, v_i\}$ by the line $\ell(wv_j)$; however this means that $\{w, v_j, z\}$ and $\{w, x, v_i\}$ form an S_1 configuration. Even more, if $x \in P \setminus V(F)$, then $\{w, x\} \notin G$ implies that this can happen on at most one side of $\ell(wx)$. We get for such an x that $|\{\{w, v_i\} : \{w, v_i, x\} \in \mathcal{T}\}| \leq 1$, hence $|\{\{w, v_i, x\} : \{w, v_i, x\} \in \mathcal{T}, 1 \leq i \leq s, x \notin V(F)\}| \leq n - 1 - s$. To estimate $|\mathcal{T}(F)|$ it remains to count the triangles from \mathcal{T} of the form $\{w, v_i, v_j\}$. For any given i there are at most two such triangles, and each of them is counted that way exactly twice, so their number is at most s . \square

Proof of Theorem 7. Suppose that $\text{del}(E(G)) \geq \lfloor n/2 \rfloor \lfloor (n - 2)/2 \rfloor$. Let the (nontrivial) components of G be F_1, F_2, \dots, F_r . Claims 4.1 and 4.2 imply that $\text{del}(E(G)) = \sum \text{del}(E(F_i)) \leq r \lfloor (n - 1)/2 \rfloor$. For n even this leads to $r \geq n/2$; equality holds, G is a perfect matching. For $n \geq 5$ odd we get $r \geq (n - 3)/2$ and in case of $r = (n - 3)/2$ we have $\text{del}(E(F_i)) = (n - 1)/2$ for each $1 \leq i \leq r$. In this latter case again Claim 4.1 implies

that each F_i has at least 4 vertices, $r \leq n/4$, a contradiction for $n > 5$. So in the odd case (for $n > 5$) we must have $r = (n - 1)/2$, each component is a single edge except perhaps one is a two-path. Then Claim 4.1 implies that $\text{del}(E(G)) \leq r \lfloor (n - 2)/2 \rfloor$, completing the proof. \square

5. Proof of Theorems 2 and 3: two separated triangles, M_1

5.1. Proof of Theorem 2 for M_1

We use a method similar to that in [15] to determine $\text{ex}_{\odot}(n, M_1)$. We prove that if H is an n -vertex M_1 -free cgh with $|H| \geq \triangle(n) + n(n - 3)/2$, then $H \in \mathcal{H}^{++}(n)$. First let $n \geq 3$ be odd. If $H \in \mathcal{H}^{++}(n)$ then we are done, so we may assume H contains a triangle $T(i, j, k) = \{v_i, v_j, v_k\}$ with $v_i < v_j < v_k < v_{i+(n-1)/2} < v_i$. Moreover, we may assume that among all such triangles, $T(i, j, k)$ is the triangle where the longest edge $\{v_i, v_k\}$ is as short as possible. Replace all triangles $T(i, j', k) \in H$ with $i < j' < k$ with all triangles $T(i - 1, k + 1, l)$ where j and l are on opposite sides of the edge $\{v_i, v_k\}$ as shown in Fig. 5. Since $T(i, j, k)$ and $T(i - 1, k + 1, l)$ form a copy of M_1 , $T(i - 1, k + 1, l) \notin H$ for all such l . Moreover, since $v_i < v_k < v_{i+(n-1)/2} < v_i$, the number of triangles $T(i - 1, k + 1, l)$ that we added is greater than the number of triangles $T(i, j, k)$ that we deleted. Consequently, this produces a cgh H' with $|H'| > |H|$. Since H is extremal M_1 -free, there exists a copy of M_1 in H' , which must contain a triangle $T(i - 1, k + 1, l) \in H'$. Since all triangles $T(i - 1, k + 1, l)$ intersect, the other triangle in the copy of M_1 must be $T(f, g, h) \in H$. Since H is M_1 -free, $T(f, g, h)$ intersects $T(i, j, k)$, which implies $v_i \leq v_f < v_g < v_h \leq v_k < v_i$ and $\{v_f, v_h\} \neq \{v_i, v_k\}$. However, then the edge $\{v_f, v_h\}$ is shorter than the edge $\{v_i, v_k\}$, a contradiction.




Fig. 5. Replacing triangles in an M_1 -free cgh.

Now let $n \geq 4$ be even and let H be an extremal n -vertex M_1 -free cgh. If $H \in \mathcal{H}^{++}(n)$ we are done, so suppose $H \notin \mathcal{H}^{++}(n)$. If H contains a triangle $T(i, j, k)$ where $v_i < v_j < v_k < v_{i+n/2-1} < v_i$, then we repeat the same proof as in the case n is odd to derive a contradiction. Therefore all triangles in H contain the centroid or are $T(i, j, k)$ with

$v_i < v_j < v_k = v_{i+n/2-1} < v_i$. The pairs $\{v_i, v_{i+n/2-1}\}$ for which there exists such a triangle $T(i, j, k)$ must pairwise intersect (possibly at their endpoints) otherwise we find a copy of M_1 in H . In particular, by Construction 3, $H \in \mathcal{H}^{++}(n)$. \square

Let us note that we can give another proof using the D_1 -graph and Theorem 2 for D_1 just as we did in subsection 4.1 to prove Theorem 2 for S_1 -free convex triangle systems – specifically, the D_1 -graph does not contain two geometrically disjoint edges.

5.2. Proof of Theorem 3 for M_1

In this section, we will prove Theorem 3 for M_1 in a stronger form using a result on geometric graphs due to Valtr [38]. A *geometric graph* (V, E) is a graph drawn in the plane so that the vertex set V consists of points in general position and the edge set E consists of straight-line segments between points of V . Two edges of a geometric graph are said to be *avoiding*, if they are opposite sides of a convex quadrilateral.

Theorem 8 (Valtr [38]). *There is a constant $C_V > 0$ such that any geometric graph on m vertices with no three pairwise avoiding edges has at most $C_V m$ edges.*

Theorem 9. *Let $n \geq 3$, C_V as above, and let \mathcal{T} be an n -point triangle system. If \mathcal{T} is M_1 -free then there exists a subfamily $\mathcal{T}' \subset \mathcal{T}$ which is D_1 -free and*

$$|\mathcal{T}| \leq |\mathcal{T}'| + C_V \binom{n}{2}.$$

Since $|\mathcal{T}'| \leq \Delta(n)$ by Theorem 3 for D_1 , one obtains the desired upper bound $|\mathcal{T}| \leq \Delta(n) + O(n^2)$.

Recall that a segment $[ab]$ (with $a, b \in P$, $a \neq b$) is a D_1 -edge in the triangle system (P, \mathcal{T}) if there are triangles from \mathcal{T} on both sides, i.e., $\exists c^-, c^+ \in P$ such that c^- and c^+ are separated by the line $\ell(ab)$ and the triangles $\{a, b, c^-\}$ and $\{a, b, c^+\} \in \mathcal{T}$. The set of all such segments is the D_1 -graph G of \mathcal{T} . For $v \in P$ let G_v be the D_1 -link graph of \mathcal{T} , i.e., it consists of those edges e of G , $v \notin e$, which are contained in a triangle $e \cup \{v\} \in \mathcal{T}$. The vertex set of the geometric graph G_v is $P \setminus \{v\}$, and for every edge $e \in G_v$ we can choose a triangle $e \cup \{-v\} \in \mathcal{T}$ which is separated from the triangle $e \cup \{v\}$ by the line $\ell(e)$, so the third vertex of $e \cup \{-v\}$ and v lie on different sides of $\ell(e)$.

Lemma 5.1. *Let \mathcal{T} be a triangle system with point set P , and let the three segments e , f , and g of $E(G_v)$ be pairwise avoiding. Then \mathcal{T} contains M_1 .*

Proof. Given a line ℓ and a set $X \neq \emptyset$ with $X \cap \ell = \emptyset$ we denote the open half plane with boundary ℓ and containing X by $H(\ell, X)$, the other side is $H(\ell, -X)$. Suppose that \mathcal{T} contains no disjoint triangles. Since e and f are on opposite sides of a convex quadrilateral, the triangle $e \cup -f \in \mathcal{T}$ should meet $f \cup \{v\}$. This is only possible if

$v \in H(\ell(e), -f)$. Similarly, $v \in H(\ell(f), -e)$, so v is in the open wedge $H(\ell(e), -f) \cap H(\ell(f), -e)$, cf., Fig. 4. For later use denote this wedge by $B(e, f)$. Since $v \in B(e, f)$, this rules out that the lines $\ell(e), \ell(f)$ are parallel.

The line $\ell(f)$ avoids the other two segments, suppose that it separates them, i.e., $e \subset H(\ell(f), -g)$ (and $g \subset H(\ell(f), -e)$). Then $B(e, f) \subset H(\ell(f), -e)$ and $B(f, g) \subset H(\ell(f), -g) = H(\ell(f), e)$. This implies $B(e, f) \cap B(f, g) = \emptyset$, contradicting to $v \in B(e, f) \cap B(f, g) \cap B(g, e)$. Hence $\ell(e)$ is a tangent line of $R := \text{conv}(\{e, f, g\})$, so this convex hull is a hexagon.

There are two cases. If R is inscribed into the triangle T formed by the lines $\ell(e)$, $\ell(f)$, and $\ell(g)$, then each region $B(e, f)$, $B(f, g)$, and $B(g, e)$ is a digon (an infinite wedge). These are pairwise disjoint, there is no place for v . Otherwise, one edge, say e lies on a side of T and f and g lie on the other two sides of the three-sided infinite region $H(\ell(e), -T) \cap H(\ell(f), g) \cap H(\ell(g), f)$. Then $B(f, g)$ is a digon inside $H(\ell(e), T)$, and $v \in B$. Consider a triangle $e \cup \{x\} \in \mathcal{T}$ where $x \in H(\ell(e), -T)$. The two digons in $H(\ell(e), -T)$ are disjoint, so we may suppose that $x \notin (H(\ell(e), -T) \cap H(\ell(f), -T))$. Then the triangle $e \cup \{x\}$ is disjoint to $f \cup \{v\}$, completing the proof of Lemma 5.1. \square

Proof of Theorem 9. Recall that we denote the set of triangles from \mathcal{T} containing an edge $e \in F$ by $\mathcal{T}(F)$, and $\text{del}(E(F))$ is the minimum number of triangles $e \cup \{x\} \in \mathcal{T}$, $e \in F$ such that removing those triangles from \mathcal{T} we eliminate all D_1 -edges of F . Our aim is to prove that $\text{del}(E(G)) \leq C_V \binom{n}{2}$ if \mathcal{T} is M_1 -free. We will show the slightly stronger statement: $\mathcal{T}(G) \leq C_V n(n-1)$. We have $\mathcal{T}(G) \leq \sum_{v \in V} |G_v|$. By Lemma 5.1 the geometric graph G_v has no three pairwise avoiding edges. Theorem 8 gives $|G_v| \leq C_V(n-1)$. Then $\text{del}(E(G)) \leq (1/2)|\mathcal{T}(G)|$ completes the proof. \square

6. Proof of Theorem 4: crossing triangles, M_3

For the proof of Theorem 4 for M_3 , it is useful to consider ordered hypergraphs: the vertex set is $\Omega_n = \{v_0, v_1, \dots, v_{n-1}\}$ with the linear ordering $v_0 < v_1 < \dots < v_{n-1}$. Let $\text{ex}_{\rightarrow}(n, M_3)$ denote the maximum number of triangles in an ordered hypergraph not containing triangles $\{v_i, v_j, v_k\}$ and $\{v_{i'}, v_{j'}, v_{k'}\}$ with $v_i < v_{i'} < v_j < v_{j'} < v_k < v_{k'}$ – this is the ordered analog of M_3 . The following theorem implies Theorem 4 for M_3 , since $\text{ex}_{\circlearrowleft}(n, M_3) = \text{ex}_{\rightarrow}(n, M_3)$:

Theorem 10. Let $n \geq 7$. Then $\text{ex}_{\rightarrow}(n, M_3) = \binom{n}{3} - \binom{n-3}{3}$.

Proof. Let H be an M_3 -free ordered triangle system with n vertices. Let H_1 consist of all $e \in H$ with $v_0, v_1 \in e$, and let H_2 consist of all $e \in H$ with $v_0 \in e$, $v_1 \notin e$ and $(e \setminus \{v_0\}) \cup \{v_1\} \in H$. Let H_3 be obtained from $H \setminus (H_1 \cup H_2)$ by merging the vertices v_0 and v_1 . Note that H_3 is a 3-cgh with $n-1$ vertices. Clearly, $|H_1| \leq n-2$. Let $G_0 = \{\{u, v\} : \{v_0, u, v\} \in H_2\}$ be the link graph of v_0 in H_2 . If two edges of G_0 cross –

say $\{u, v\}, \{w, x\} \in G_0$ with $u < w < v < x$, then the triangles $\{u, v, v_1\}$ and $\{w, x, v_0\}$ are in H_2 , and form a copy of M_3 , a contradiction. Therefore no two edges of G_0 cross, which implies G_0 is an outerplane graph with $n - 2$ vertices. Consequently $|G_0| \leq 2n - 7$, by Euler's Formula. Finally, it is also straightforward to check H_3 is M_3 -free, so by induction,

$$|H| = |H_1| + |H_2| + |H_3| \leq (n - 2) + (2n - 7) + \binom{n-1}{3} - \binom{n-4}{3} = \binom{n}{3} - \binom{n-3}{3}.$$

This completes the proof of Theorem 10. \square

7. Proof of Theorem 4: stabbing triangles, M_2

We prove by induction on n that $\text{ex}_\triangle(n, M_2) = \binom{n}{2} - 2$ for $n \geq 7$. When $n = 7$, since cyclically consecutive triangles $\{v_i, v_{i+1}, v_{i+2}\}$ are never in M_2 , we may assume these seven triangles are in any M_2 -free cgh. For the remaining twenty-eight triangles, we create a graph with vertex set consisting of these triangles and form an edge if two of the triangles form a copy of M_2 . A computer aided calculation [35] then yields this graph has independence number 12 and hence $\text{ex}_\triangle(7, M_2) = 12 + 7 = \binom{7}{2} - 2$.

For the induction step, we plan to find two consecutive $u, v \in \Omega_n$ with degree at most three and whose common link graph $G_u \cap G_v$ has at most $n - 3$ edges. Let H be a maximal M_2 -free cgh on Ω_n , and $H' \subset H$ be the cgh after removing all consecutive triangles $\{v_i, v_{i+1}, v_{i+2}\}$. Let $d(v_i, v_j)$ be length of the path on the perimeter of the polygon starting with v_i and moving clockwise to v_j . For a triangle $e = \{v_i, v_{i+1}, v_k\} \in H'$ – we only consider such triangles – let $\ell(e) = \min\{d(v_{i+1}, v_k), d(v_k, v_i)\}$.

Lemma 7.1. *Let $H \subset \binom{\Omega_n}{3}$ be a maximal M_2 -free cgh and H' be as above. Then*

- (1) *For consecutive $u, v \in \Omega_n$, $|G_u \cap G_v| \leq n - 3$ with equality only if $G_u \cap G_v$ is a star.*
- (2) *There exists $v_i \in \Omega_n$ such that the degree of $\{v_i, v_{i+1}\}$ is at most three in H .*

Proof. We first prove (1) by showing $G_{u,v} := G_u \cap G_v$ does not contain a pair of disjoint edges. If $\{w, x\}, \{y, z\}$ are disjoint edges in $G_{u,v}$, and $v < w < x < y < z < u < v$ or $v < w < y < z < x < u < v$ – this means that $\{w, x\}, \{y, z\}$ do not cross – then $\{u, w, x\}, \{v, y, z\}$ form M_2 . If on the other hand $v < w < y < x < z < u < v$ – this means $\{w, x\}, \{y, z\}$ do cross – then $\{u, y, z\}, \{v, w, x\}$ form M_2 . So $G_{u,v}$ has no pair of disjoint edges. It is a standard fact that the unique extremal graphs with at least four vertices and no pair of disjoint edges are stars, and therefore $G_{u,v}$ has at most $n - 3$ edges.

For (2), seeking a contradiction, suppose $d_H(\{v_i, v_{i+1}\}) \geq 4$ (and hence $d_{H'}(\{v_i, v_{i+1}\}) \geq 2$) for every $v_i \in \Omega_n$. We first show there exists $e \in H'$ with $\ell(e) \geq 3$. If not, then $\{v_i, v_{i+1}, v_{i+3}\} \in H'$ and $\{v_{i-2}, v_i, v_{i+1}\} \in H'$ for all i and there are no other triangles in H' . However, then $\{v_0, v_1, v_3\} \in H'$ and $\{v_2, v_4, v_5\} \in H'$ form M_2 , a

contradiction. So there exists $e \in H'$ with $\ell(e) \geq 3$. From all $e \in H'$ with $\ell(e) \geq 3$, pick e so that $\ell(e) = j \geq 3$ is a minimum. Suppose $e = \{v_0, v_1, v_{j+1}\}$, so $\ell(e) = d(v_1, v_{j+1})$ (the proof for e of the form $\{v_{n-j}, v_0, v_1\}$ with $\ell(e) = j = d(v_{n-j}, v_0) \geq 3$ will be symmetric). Then $d_{H'}(\{v_{j-1}, v_j\}) \geq 2$, so there are triangles $f = \{v_h, v_{j-1}, v_j\}$ and $g = \{v_k, v_{j-1}, v_j\}$ in H' . If $j+1 < k \leq n-1$ or $j+1 < h \leq n-1$, then f and e or g and e respectively form M_2 , a contradiction. So $0 \leq h, k \leq j-3$, recalling $\{v_{j-2}, v_{j-1}, v_j\} \notin H'$. Now

$$\ell(f) = d(v_h, v_{j-1}) > d(v_k, v_{j-1}) \geq 2$$

and so $\ell(f) \geq 3$. On the other hand, since $0 \leq h < j-1$,

$$\ell(f) = d(v_h, v_{j-1}) < d(v_0, v_j) = \ell(e)$$

contradicting the choice of e . This final contradiction proves (2). \square

Let $\{v_i, v_{i+1}\}$ have degree at most three in H , as guaranteed by Lemma 7.1 part (2). We contract the edge $\{v_i, v_{i+1}\}$ to a vertex w to get a cgh H_0 with $n-1$ vertices. Let $G = \{\{u, v\} : \{u, v, v_i\}, \{u, v, v_{i+1}\} \in H\}$ be the common link graph of v_i and v_{i+1} .

Lemma 7.2. *Let G be the common link graph of v_i and v_{i+1} . Then $|G| \leq n-4$.*

Proof. If neither of $\{v_{i-1}, v_i, v_{i+2}\}$ or $\{v_{i-1}, v_{i+1}, v_{i+2}\}$ is in H , then $\{v_{i-1}, w, v_{i+2}\} \notin H_0$ and $|G| \leq n-4$ follows from Lemma 7.1 part (1). So we assume $\{v_{i-1}, v_i, v_{i+2}\} \in H$ or $\{v_{i-1}, v_{i+1}, v_{i+2}\} \in H$.

Case 1. $\{v_{i-1}, v_i, v_{i+2}\} \in H$. Suppose G is a star with $n-3$ edges, with center v_k . If $v_k \notin \{v_{i-1}, v_{i+2}\}$, then letting $v_j \notin \{v_k, v_{i-1}, v_i, v_{i+1}, v_{i+2}\}$, it follows that $\{v_i, v_j, v_k\}$ and $\{v_{i-1}, v_{i+1}, v_{i+2}\}$ form a copy of M_2 . Hence, we may assume that $v_k = v_{i-1}$ or $v_k = v_{i+2}$. Both of these cases are similar, so consider only the case $v_k = v_{i+2}$. We may assume that $\{v_{i+3}, v_{i+4}\}$ has degree at least three. Then there is at least one triangle which contains $\{v_{i+3}, v_{i+4}\}$ of the form $\{v, v_{i+3}, v_{i+4}\}$. If $v \in \Omega_n$ and $v_{i+4} < v < v_{i+1} < v_{i+3}$, then $\{v, v_{i+3}, v_{i+4}\}$ and $\{v_{i+1}, v_{i+2}, v_{i+5}\}$ form M_2 . If $v = v_{i+1}$, then $\{v, v_{i+3}, v_{i+4}\}$ and $\{v_{i-1}, v_i, v_{i+2}\}$ form M_2 . So G is not a star with $n-3$ edges, and Lemma 7.1 part (1) gives $|G| \leq n-4$.

Case 2. $\{v_{i-1}, v_{i+1}, v_{i+2}\} \in H$. In this case, a symmetric argument to that used for $\{v_{i-1}, v_i, v_{i+2}\} \in H$ applies by reversing the orientation of Ω_n . \square

To complete the proof of $|H| \leq \binom{n}{2} - 2$, we note by inspection that H_0 is also M_2 -free. By induction, $|H_0| \leq \binom{n-1}{2} - 2$. By Lemma 7.2, and recalling $d_H(v_i, v_{i+1}) \leq 3$,

$$|H| = |H_0| + |G| + d_H(v_i, v_{i+1}) \leq \binom{n-1}{2} - 2 + n - 4 + 3 = \binom{n}{2} - 2.$$

This proves Theorem 4 for M_2 . \square

8. Proof of Theorem 4: crossing triangles sharing a vertex, S_3

Let $H \subset \binom{\Omega_n}{3}$ be a S_3 -free cgh and G_i be the link graph of v_i in H . Let G'_i comprise the edges of G_i which consist of two consecutive vertices in Ω_n , and let $G''_i = G_i \setminus G'_i$.

Lemma 8.1. *Let $H \subset \binom{\Omega_n}{3}$ be a S_3 -free cgh. For $0 \leq i \leq n-1$, $|G''_i| \leq n-3$.*

Proof. The graph G''_i has no pair of crossing edges since H is S_3 -free. If we add to G''_i all the n edges $\{v_j, v_{j+1}\}$, we obtain an outerplanar graph which has at most $2n-3$ edges. Removing the n added edges gives $|G''_i| \leq n-3$. \square

Lemma 8.2. *Let $H \subset \binom{\Omega_n}{3}$ be a S_3 -free cgh. For each i , $|G'_i| + |G'_{i+1}| \leq n$.*

Proof. We may assume $i = 0$. Let G denote the multigraph obtained by superimposing the graphs G'_0 and G'_1 , so $|G| = |G'_0| + |G'_1|$. As G'_i comprises the edges of G_i which consist of two consecutive vertices in Ω_n , each component C of G is a path P with some edges of multiplicity two. If $\{v_{j-1}, v_j\} \in P \cap G'_0$, then $\{v_j, v_{j+1}\} \notin P \cap G'_1$, otherwise $\{v_0, v_j, v_{j+1}\}, \{v_1, v_{j-1}, v_j\}$ form $S_3 \subset H$ as in Fig. 6, a contradiction. If all edges of P are from G'_1 only, then $|C| = |P| = |V(C)| - 1$. Otherwise, let $\{v_j, v_{j+1}\}$ be the first edge of P in G'_0 in the clockwise direction. Then all edges of P preceding $\{v_j, v_{j+1}\}$ are in G'_1 only, and all edges of P after $\{v_j, v_{j+1}\}$ are in G'_0 only, whereas $\{v_j, v_{j+1}\}$ might be in both G'_0 and in G'_1 . Therefore at most one edge of P has multiplicity two, and $|C| \leq |P| + 1 = |V(C)|$. If C_1, C_2, \dots, C_r are the components of G , we conclude $|G| = |C_1| + |C_2| + \dots + |C_r| \leq |V(C_1)| + |V(C_2)| + \dots + |V(C_r)| = |V(G)| = n$. \square




Fig. 6. Crossing triangles in the proof of Lemma 8.2.

We now complete the proof of $\text{ex}_\cup(n, S_3) \leq n(n-2)/2$, using the following identity:

$$3|H| = \sum_i (|G'_i| + |G''_i|) = \sum_i \frac{1}{2} (|G'_i| + |G'_{i+1}|) + \sum_i |G''_i|.$$

We apply Lemmas 8.1 and 8.2 to each term in the sums to obtain:

$$3|H| \leq \sum_{i=0}^{n-1} \frac{1}{2}n + \sum_{i=0}^{n-1} (n-3) = \frac{1}{2}n^2 + n(n-3) = \frac{3}{2}n(n-2). \quad \square$$

9. Proof of Theorem 5: touching triangles with parallel sides, S_2

Let $H \subset \binom{\Omega_n}{3}$ be an S_2 -free cgh. We are going to show $|H| \leq 23n^2/64$. Consider a triangle $e = \{v_i, v_j, v_k\} \in H$ where $v_i < v_j < v_k < v_i$. We call the edge $\{v_i, v_j\}$ *good for* e if there does not exist a k' such that $v_j < v_{k'} < v_k < v_j$ and $\{v_i, v_j, v_{k'}\} \in H$, and *bad* otherwise.

Lemma 9.1. *Let $H \subset \binom{\Omega_n}{3}$ be an S_2 -free cgh. Then*

- (1) *Every triangle of H contains at least two good edges.*
- (2) *Every edge in ∂H is good for either one or two triangles of H .*

Proof. We first prove (1). Suppose $e = \{v_i, v_j, v_k\} \in H$ and $\{v_i, v_j\}$ and $\{v_j, v_k\}$ are bad. Then there exist $k' : v_j < v_{k'} < v_k < v_j$ and $i' : v_k < v_{i'} < v_i < v_k$ such that $\{v_i, v_j, v_{k'}\}, \{v_j, v_k, v_{i'}\} \in H$. However, the triangles $\{v_{i'}, v_j, v_k\}$ and $\{v_i, v_j, v_{k'}\}$ form configuration S_2 , a contradiction.

For (2), given $\{v_i, v_j\} \in \partial H$, consider a triangle $\{v_i, v_j, v_k\}$ with $v_i < v_j < v_k < v_i$ and v_k as close as possible to v_j ; this determines v_k uniquely. Similarly, for $\{v_i, v_j\} \in \partial H$, consider a triangle $\{v_i, v_j, v_k\}$ with $v_i < v_k < v_j < v_i$ with v_k as close as possible to v_i ; this too determines v_k uniquely. Therefore each edge in ∂H is good for either one of two triangles of H . \square

Color an edge in ∂H *blue* if it is good for exactly one triangle in H , and *red* if it is good for exactly two triangles in H . Let R be the number of red edges and B the number of blue edges – for a red edge $\{u, v\}$, there exist vertices $w, x \in \Omega_n$ on opposite sides of $\{u, v\}$ such that $\{u, v, w\} \in H$ and $\{u, v, x\} \in H$, so red edges are what we have referred to as edges in the D_1 -graph in this paper. If we map a triangle $e \in H$ to the edges in e that are good for e , then each red edge is counted twice and each blue edge is counted once. On the other hand, each triangle of H contains at least two good edges, by Lemma 9.1, so $2|H| \leq 2R + B$. In particular,

$$|H| \leq R + B/2 \leq R + B = |\partial H|.$$

Lemma 9.2. *If $\{v_i, v_j\}$, $\{v_j, v_k\}$ and $\{v_k, v_i\}$ are red edges, then $\{v_i, v_j, v_k\} \in H$.*

Proof. Suppose $\{v_i, v_j, v_k\} \notin H$ and $v_i < v_j < v_k < v_i$. Then by definition there exists $k' \neq k$ such that $\{v_i, v_j, v_{k'}\} \in H$ and $v_j < v_{k'} < v_i < v_j$. We consider two cases.

Case 1. $v_j < v_{k'} < v_k < v_j$. There exists $i' \neq i$ such that $\{v_{i'}, v_j, v_k\} \in H$ and $v_k < v_{i'} < v_j < v_k$. We observe $v_i < v_{i'} < v_j < v_i$, otherwise $\{v_j, v_{k'}\}$ and $\{v_i, v_{i'}\}$ are non-crossing, and $\{v_i, v_j, v_{k'}\}$ and $\{v_{i'}, v_j, v_k\}$ form S_2 in H . Now there exists $j' \neq j$

such that $\{v_i, v_{j'}, v_k\} \in H$ and $v_i < v_{j'} < v_k < v_i$. If $v_i < v_{j'} < v_j < v_i$, then the edges $\{v_{j'}, v_k\}$ and $\{v_j, v_{k'}\}$ are non-crossing, and $\{v_i, v_j, v_{k'}\}$ and $\{v_i, v_j, v_k\}$ form S_2 . If $v_j < v_{j'} < v_k < v_j$, then $\{v_{i'}, v_j\}$ and $\{v_i, v_j\}$ are “parallel”, and $\{v_{i'}, v_j, v_k\}$ and $\{v_i, v_j, v_{k'}\}$ form S_2 in H .

Case 2. $v_k < v_{k'} < v_i < v_k$. Consider the reverse ordering of Ω_n and apply the proof of Case 1. \square

By Lemma 9.2, every triangle of red edges is a triangle of H , so there are at most $|H| \leq |\partial H| \leq \binom{n}{2}$ such triangles. In particular, the number of red edges is at most $n^2/4 + n/2$ – one could use a precise result by Lovász-Simonovits [27] to deduce this. Instead we give a direct proof: the number of triangles in any graph G is at least

$$\sum_{\{u,v\} \in E(G)} (d(u) + d(v) - n).$$

If G has average degree d , then this is precisely

$$\sum_u d(u)^2 - \frac{1}{2}dn^2 \geq d^2n - \frac{1}{2}dn^2.$$

Since the graph G of red edges in ∂H has at most $|H| \leq \binom{n}{2}$ triangles,

$$d^2n - \frac{1}{2}dn^2 \leq \frac{1}{2}n^2$$

which gives $d \leq n/2 + 1$ and therefore $R = |G| \leq n^2/4 + n/2$. Therefore

$$2|H| \leq 2R + B \leq \binom{n}{2} + \left(\frac{n^2}{4} + \frac{n}{2}\right) = \frac{3n^2}{4}.$$

To improve this bound to the desired $|H| \leq 23n^2/64$, we may assume n is odd and partition the complete graph on Ω_n into planar matchings M_1, M_2, \dots, M_n where $M_i = \{\{v_j, v_k\} : j+k \equiv i \pmod n\}$. Then there exists $i \leq n$ such that at least R/n edges in $M = M_i$ are red. For each pair of red edges, say $\{u, v\}$ and $\{w, x\}$, where $u < w < x < v < u$, there exist triangles $\{u, v, y\}, \{w, x, z\} \in H$ where $u < w < z < x < v < y < u$. Now by inspection, the edge $\{y, z\}$ cannot be contained in any triangle of H without creating configuration S_2 – see Fig. 7. Furthermore, if $\{u', v'\}, \{w', x'\} \in M$, then $\{u', v', y\}$ and $\{w', x', z\}$ cannot both be triangles of H without creating S_2 . Therefore for each pair $\{\{u, v\}, \{w, x\}\}$ of red edges of M , we may associate a unique edge $\{y, z\}$ which is not contained in any triangle of H . Consequently

$$2|H| \leq 2R + B \leq 2R + \binom{n}{2} - \binom{R/n}{2} - R \leq R + \binom{n}{2} - \binom{R/n}{2}.$$

Since $R \leq n^2/4 + n/2$, this implies $|H| \leq 23n^2/64 - n/4 + 3/8$. As $n \geq 3$, this is at most $23n^2/64$, as required. \square




Fig. 7. Edge $\{y, z\}$ absent from ∂H . (For interpretation of the colors in the figure(s), the reader is referred to the web version of this article.)

10. Proof of Theorem 6: triangles sharing a side, D_2

We first observe some simple bounds on $\text{ex}_\circ(n, D_2)$. If G is a convex geometric graph that is a triangulation of a convex polygon, then the family $T(G)$ of vertex sets of the triangular regions in G form a D_2 -free cgh. By Euler's Formula, $|T(G)| < \frac{1}{2}|G|$, so if G_1, G_2, \dots, G_M are edge-disjoint triangulations of polygons with vertices from Ω_n , then $H = T(G_1) \cup T(G_2) \cup \dots \cup T(G_M)$ is a D_2 -free cgh on Ω_n . Each D_2 -free cgh H can be obtained in this way, so we get $\text{ex}_\circ(n, D_2) < (1/2)\binom{n}{2}$. On the other hand, every Steiner triple system induces a D_2 -free cgh, we get $\text{ex}_\circ(n, D_2) \geq \frac{1}{3}\binom{n}{2} - O(n)$. Construction 8 improves this to $\frac{3}{7}\binom{n}{2} - O(n)$, and Damásdi and N. Frankl [11] showed $\text{ex}_\circ(n, D_2) \geq \frac{2n^2-3n}{9}$ for all $n \equiv 6 \pmod{9}$ by a different method. Here we prove the upper bound $\text{ex}_\circ(n, D_2) \leq \frac{2n^2-3n}{9}$ for all n .

For the calculation below we need a simple proposition which can be shown by standard high school calculus. If $h, x \geq 0$ are reals, $n \geq 3$ is an integer and $h \geq (2n-3)/9$, then

$$(h+2x)(h+2x+1) \leq 2xn \implies x \geq \frac{n+3}{18}. \quad (4)$$

Another elementary proposition is the following statement: Suppose that A is a *multiset* of positive integers such that the multiplicity of each entry is at most n , then

$$\sum_{a \in A} a \geq \frac{|A|(|A|+n)}{2n}. \quad (5)$$

For the upper bound on $\text{ex}_\circ(n, D_2)$, let $H \subset \binom{\Omega_n}{3}$ be a D_2 -free cgh. The graph ∂H has a (unique) edge-disjoint decomposition into triangulations G_1, \dots, G_M as follows. Make a graph C with vertex set H : two triangles of H are joined by an edge of C if they

share a side. Consider the partition of C generated by the components C_1, C_2, \dots, C_M of G , where $|C_i| = k_i$. Each C_i corresponds to a hypergraph $H_i \subset H$ of triangles. Since H_i is D_2 -free each $G_i := \partial H_i$ forms a triangulation of a convex $(k_i + 2)$ -gon P_i with $k_i - 1$ diagonals, $T(G_i) = H_i$, $|E(G_i)| = 2k_i + 1$. Let A_i be the multiset of integers consisting of the side lengths of P_i , $|A_i| = k_i + 2$. We have

$$\sum_{a \in A_i} a \leq n \quad (6)$$

and here equality holds if the polygon P_i contains the center of Ω_n . Let A be the multiset $\cup_{i \leq M} A_i$. Since each edge of ∂H appears in exactly one G_i and there are n (or $n/2$ or 0) diagonals of Ω_n of a given length we obtain that A is a multiset with maximum multiplicities at most n . Moreover, $|A| = \sum_i (k_i + 2) = |H| + 2M$, so (5) and (6) yield

$$\frac{(|H| + 2M)(|H| + 2M + n)}{2n} \leq \sum_{a \in A} a = \sum_{i \leq M} \left(\sum_{a \in A_i} a \right) \leq Mn. \quad (7)$$

Suppose that $|H| \geq (2n^2 - 3n)/9$. Define h, x as $h := |H|/n$ and $x := M/n$. Then $h \geq (2n - 3)/9$ and (7) and (4) imply $x \geq (n + 3)/18$. However

$$2|H| + M = \sum_{1 \leq i \leq M} (2k_i + 1) = \sum |E(G_i)| = |\partial H| \leq \binom{n}{2}.$$

Hence $|H| \leq \frac{1}{2}(\binom{n}{2} - xn) \leq (2n^2 - 3n)/9$. \square

11. Concluding remarks

- In this paper, we considered convex geometric configurations consisting of two triangles. One may consider analogous problems for r -tuples: for instance, how many r -gons can a convex geometric n -vertex r -graph have if it does not contain two r -gons which are geometrically disjoint (this is the r -uniform analog of M_1)? This problem was posed explicitly by P. Frankl, Holmsen and Kupavskii [15]:

Problem 11.1. *Find analogues of our results for other classes of sets such as convex r -gons in \mathbb{R}^2 .*

A family of convex r -gons in the plane is *strongly intersecting* if any two of the members share a point in their interior. The maximum size of a strongly intersecting family of r -gons is obtained from the obvious extensions of Construction 1. Consider the family of all r -gons containing the centroid of Ω_n when n is odd, together with, for each diameter ℓ , all r -gons which have a side equal to ℓ and which lie on one side of ℓ . Letting $\triangle_r(n)$ denote the size of these families, it is not hard to see

$$\Delta_r(n) = \binom{n}{r} - n \binom{(n-1)/2}{r-1}$$

if n is odd, and $\Delta_r(n)$ can be computed similarly if n is even. In particular, $\Delta_r(n) = (1 - r/2^{r-1})\binom{n}{r} + O(n^{r-1})$ for each $r \geq 3$.

Theorem 11. *The maximum size of a strongly intersecting family of r -gons from Ω_n is $\Delta_r(n)$.*

Proof. (Sketch). We proceed in a similar way to the proof of Theorem 2 for M_1 . Consider any r -gon $\{v_{i_1}, v_{i_2}, \dots, v_{i_r}\}$ in H with $v_{i_1} < v_{i_2} < \dots < v_{i_r} < v_{i_1}$ and where the longest side $\{v_{i_1}, v_{i_r}\}$ is as short as possible, and replace all such r -gons with $\{v_{i_1}, v_{j_2}, \dots, v_{j_{r-1}}, v_{i_r}\}$ where $v_{i_r} < v_{j_2} < v_{j_3} < \dots < v_{j_{r-1}} < v_{i_1}$. Since the number of choices of j_2, j_3, \dots, j_{r-1} is always at least the number of choices of i_2, i_3, \dots, i_{r-1} , this new r -cgh H' has $|H'| \geq |H|$. So we repeat until H' consists of all r -gons containing the centroid of Ω_n when n is odd, or n is even and H' consists of all r -gons containing the centroid plus for each diameter ℓ all r -gons which have a side equal to ℓ and which lie on one side of ℓ . \square

- Since there are many other possible configurations of two r -gons, we did not discuss these problems in this paper. Some special cases were studied in [16]: for instance, if F consists of two r -gons $\{u_1, u_2, \dots, u_r\}$ and $\{v_1, v_2, \dots, v_r\}$ where $u_1 < v_1 < u_2 < v_2 < \dots < u_r < v_r < u_1$, then it was shown in [16] that for $n > r > 1$,

$$\text{ex}_{\cup}(n, F) = \binom{n}{r} - \binom{n-r}{r}.$$

This may be viewed as a geometric or ordered version of the Erdős-Ko-Rado Theorem [12].

- In the cases of M_2, M_3 and S_3 (see Fig. 1), we obtained exact results for the extremal functions in convex geometric hypergraphs / convex triangle systems (for n even in the case of S_3). Our proofs, with more work, should give a characterization of the extremal examples as well. For M_2 , one requires $n \geq 8$ for the extremal configuration to be unique, as verified by computer. For S_3 , we believe that $\text{ex}_{\cup}(n, S_3) = (n-1)(n-2)/2 + 1$ when n is odd, but do not have a proof, and we also do not know the characterization of extremal S_3 -free convex triangle systems (this is the content of Problem 1.3).

- It is likely the case that most of our theorems hold equally for *ordered hypergraphs*, where the vertex set is linearly ordered, but we did not work out the details except for the obvious case M_3 (see the first paragraph in Section 6). The case of S_2 stands out, since the ordered extremal number is not the same as the convex geometric extremal number. The ordered construction would be to take all triangles $\{v_i, v_{i+1}, v_j\}$ from an ordered vertex set $\{v_0, v_1, \dots, v_{n-1}\}$ where $i \geq 0$ and $i+1 < j \leq n-1$.

Extremal problems for matchings in ordered graphs connect to enumeration of permutations [28] and these have also been extended to hypergraphs [24].

- A hypergraph H is *linear* if for distinct hyperedges $e, f \in E(H)$, $|e \cap f| \leq 1$. The extremal functions for the configurations in this paper in the context of linear cghs were determined in [2] up to constant factors for all the configurations except S_2 . Specifically, if $\text{ex}_\odot^*(n, F)$ is the maximum number of triangles in an n -vertex F -free linear cgh, then Aronov, Dujmović, Morin, Ooms and da Silveira [2] proved $\text{ex}_\odot^*(n, M_2) = \Theta(n)$, whereas if $F \in \{M_1, M_3, S_1, S_3\}$, $\text{ex}_\odot^*(n, F) = \Theta(n^2)$. It would be interesting to determine the exact extremal functions in each case. The problem of determining $\text{ex}_\odot^*(n, S_2)$ appears to be very difficult, as it is connected to monotone matrices, tripod packing, and 2-comparable sets – see Aronov, Dujmović, Morin, Ooms and da Silveira [2] for details. The best bounds are $\text{ex}_\odot^*(n, S_2) = \Omega(n^{1.546})$ due to Gowers and Long [19] and $\text{ex}_\odot^*(n, S_2) = n^2 / \exp(\Omega(\log^* n))$ due to the best bounds on the removal lemma by Fox [14].

- By a result of Boros and Füredi [5], for every n -point set P (no three on a line) one can find a point on the plane which is contained in at least $n^3/27 - O(n^2)$ triangles with these vertices; and Bukh, Matoušek, and Nivasch [9] gave an example that the coefficient $1/27$ is the best possible. It would be interesting to determine the largest subsystem of pairwise intersecting triangles in this construction.

- One can further relax the conditions on the point sets to allow *all* planar n -point sets. We conjecture that our upper bounds in Theorem 3 hold for all planar n -point sets (when we only count the proper triangles with non-empty interiors). Surely in that case one has to relax the definition of configurations (like, e.g., Ackerman, Nitzan, and Pinchasi [1] did about avoiding pairs of edges).

- We have not considered F -free triangle systems (P, \mathcal{T}) where the point set P is not necessarily in convex position and $F \in \{M_2, M_3, S_2, S_3, D_2\}$. The reason is, unlike in the case $F \in \{D_1, S_1, M_1\}$, there are many different ways to extend the definitions of these configurations and these can lead to many different problems. E.g., if one insists that no triangle in F contains another vertex of F then the answer is always at least $n^3/27 + O(n^2)$ as it is shown by the following example $P := X \cup Y \cup Z$, $\mathcal{T} := \{xyz : x \in X, y \in Y, z \in Z\}$ and $X := \{(i, 10^{-i}) : 1 \leq i \leq n/3\}$, $Y := \{(10^{-i}, i) : 1 \leq i \leq n/3\}$, and $Z := \{(-i, -i + 10^{-i}) : 1 \leq i \leq n/3\}$. It is a rich area with full of problems, e.g., it would be interesting to determine all configurations F satisfying that $|\mathcal{T}| \leq (1 + o(1))\text{ex}_\odot(n, F)$ holds for F -free triangle systems.

Acknowledgments

The research of the first author was supported in part by NKFIH grant KH130371 and NKFI-133819. The second author was supported by NSF award DMS-1952767 and the last two authors were supported by NSF award DMS-1800332. The authors are very grateful to both referees for carefully reading the paper and providing many valuable comments that improved the presentation.

References

- [1] E. Ackerman, N. Nitzan, R. Pinchasi, The maximum number of edges in geometric graphs with pairwise virtually avoiding edges, *Graphs Comb.* 30 (2014) 1065–1072.
- [2] B. Aranov, V. Dujmvić, P. Morin, A. Ooms, L. Xavier da Silveira, More Turán-type theorems for triangles in convex point sets, *Electron. J. Comb.* 26 (2019).
- [3] I. Bárány, A generalization of Carathéodory’s theorem, *Discrete Math.* 40 (1982) 141–152.
- [4] E. Boros, Z. Füredi, Su un teorema di Kármán nella geometria combinatoria, *Archimedes* 29 (1977) 71–76 (in Italian).
- [5] E. Boros, Z. Füredi, The number of triangles covering the center of an n -set, *Geom. Dedic.* 17 (1984) 69–77.
- [6] P. Braß, Turán-type extremal problems for cghs, in: *Towards a Theory of Geometric Graphs*, in: *Contemp. Math.*, vol. 342, Amer. Math. Soc., Providence, RI, 2004, pp. 25–33.
- [7] P. Braß, G. Rote, K.J. Swanepoel, Triangles of extremal area or perimeter in a finite planar point set, *Discrete Comput. Geom.* 26 (2001) 51–58.
- [8] B. Bukh, A point in many triangles, *Electron. J. Comb.* 13 (2006).
- [9] B. Bukh, J. Matoušek, G. Nivasch, Stabbing simplices by points and flats, *Discrete Comput. Geom.* 43 (2010) 321–338.
- [10] V. Capovelas, J. Pach, A Turán-type theorem for chords of a convex polygon, *J. Comb. Theory, Ser. B* 56 (1992) 9–15.
- [11] G. Damásdi, N. Frankl, D_2 -free cghs, manuscript, December 2020.
- [12] P. Erdős, C. Ko, R. Rado, Intersection theorems for systems of finite sets, *Q. J. Math.* 12 (1961) 313–320.
- [13] P. Erdős, J. Spencer, *Probabilistic Methods in Combinatorics, Probability and Mathematical Statistics*, vol. 17, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1974, 106 pp.
- [14] J. Fox, A new proof of the graph removal lemma, *Ann. Math.* 174 (2011) 561–579.
- [15] P. Frankl, A. Holmsen, A. Kupavskii, Intersection theorems for triangles, <https://arxiv.org/abs/2009.14560>, 2020.
- [16] Z. Füredi, T. Jiang, A. Kostochka, D. Mubayi, J. Verstraete, Extremal problems on ordered and convex geometric hypergraphs, *Can. J. Math.* (2020) 1–21.
- [17] Z. Füredi, T. Jiang, A. Kostochka, D. Mubayi, J. Verstraete, Tight paths in convex geometric hypergraphs, *Adv. Comb.* (2020).
- [18] Z. Füredi, T. Jiang, A. Kostochka, D. Mubayi, J. Verstraete, Ordered and convex geometric trees with linear extremal function, *Discrete Comput. Geom.* 64 (2020) 324–338.
- [19] W.T. Gowers, J. Long, The length of an s -increasing sequence of r -tuples, <https://arxiv.org/abs/1609.08688>, 2016.
- [20] M. Gromov, Singularities, expanders and topology of maps. Part 2: from combinatorics to topology via algebraic isoperimetry, *Geom. Funct. Anal.* 20 (2010) 416–526.
- [21] H. Hopf, E. Pannwitz, Aufgabe Nr. 167, *Jahresber. Dtsch. Math.-Ver.* 43 (1934) 114.
- [22] R. Karasëv, A simpler proof of the Boros-Füredi-Bárány-Pach-Gromov theorem, *Discrete Comput. Geom.* 47 (2012) 492–495.
- [23] C. Keller, M. Perles, On convex geometric graphs with no $k + 1$ pairwise disjoint edges, *Graphs Comb.* 32 (2016) 2497–2514.
- [24] M. Klazar, A. Marcus, Extensions of the linear bound in the Füredi-Hajnal conjecture, *Adv. Appl. Math.* 38 (2007) 258–266.
- [25] Y.S. Kupitz, On pairs of disjoint segments in convex position in the plane, *Ann. Discrete Math.* 20 (1984) 203–208.
- [26] Y.S. Kupitz, M. Perles, Extremal theory for convex matchings in convex geometric graphs, *Discrete Comput. Geom.* 15 (1996) 195–220.
- [27] L. Lovász, M. Simonovits, On the number of complete subgraphs of a graph II, in: *Studies in Pure Mathematics*, 1983, pp. 459–495.
- [28] A. Marcus, G. Tardos, Excluded permutation matrices and the Stanley-Wilf conjecture, *J. Comb. Theory, Ser. A* 107 (2004) 153–160.
- [29] J.W. Moon, *Topics on Tournaments*, Holt, Rinehart and Winston, New York-Montreal, Que.-London, 1968, viii+104 pp.
- [30] J. Pach, Geometric graph theory, in: *Surveys in Combinatorics*, 1999, Canterbury, in: *London Math. Soc. Lecture Note Ser.*, vol. 267, Cambridge Univ. Press, Cambridge, 1999, pp. 167–200.

- [31] J. Pach, The beginnings of geometric graph theory, in: Erdős Centennial, in: Bolyai Soc. Math. Stud., vol. 25, János Bolyai Math. Soc., Budapest, 2013, pp. 465–484.
- [32] J. Pach, R. Pinchasi, How many unit equilateral triangles can be generated by n points in general position?, *Am. Math. Mon.* 110 (2003) 100–106.
- [33] J. Pach, R. Pinchasi, G. Tardos, G. Tóth, Geometric graphs with no self-intersecting path of length three, *Eur. J. Comb.* 25 (2004) 793–811.
- [34] J. Pach, G. Tardos, Forbidden paths and cycles in ordered graphs and matrices, *Isr. J. Math.* 155 (2006) 359–380.
- [35] The Sage Developers, SageMath, the Sage Mathematics software system (Version 9.1), <https://www.sagemath.org/>, 2020.
- [36] J.W. Sutherland, Lösung der Aufgabe 167, *Jahresber. Dtsch. Math.-Ver.* 45 (1935) 33–35.
- [37] G. Tardos, Extremal theory of ordered graphs, in: *Proceedings of the International Congress of Mathematics – 2018*, vol. 3, 2019, pp. 3219–3228.
- [38] P. Valtr, On geometric graphs with no k pairwise parallel edges, *Discrete Comput. Geom.* 19 (1998) 461–469.
- [39] U. Wagner, E. Welzl, A continuous analogue of the upper bound theorem, *Discrete Comput. Geom.* 26 (2001) 205–219.
- [40] R.M. Wilson, An existence theory for pairwise balanced designs, I and II, *J. Comb. Theory, Ser. A* 13 (1972) 220–245 and 246–273.