

CORRIGENDUM

Rational approximation on the positive real axis

[*Proc. London Math. Soc.* (3) 31 (1975) 439-56]

By PAUL ERDÖS and A. R. REDDY

[Received 17 November 1976]

There is an inaccuracy in one of the calculations presented in the proof of Theorem 2 (pp. 443-44). This occurred in equation (20).

By making the following changes throughout the proof:

$$\begin{array}{ll} D(n) & \text{to } [D(n)]^{\frac{1}{2}}, \\ e_{k+1}(x) & \text{to } e_2(x), \\ l_k(n[D(n)]^{-1}) & \text{to } l_1(n[D(n)]^{-1}), \\ l_k(ne^2[D(n)]^{-1}) & \text{to } l_1(ne^2[D(n)]^{-1}), \end{array}$$

we get easily the following:

THEOREM A. *For every $c > 0$ and $k \geq 1$ there is an entire function of infinite order with non-negative coefficients for which there exist infinitely many n such that*

$$\lambda_{0,n} \geq \exp(-(n[D(n)]^{-1})).$$

By careful calculation $\frac{1}{2}$ can be replaced by $\frac{1}{2} - \varepsilon$ ($\varepsilon > 0$).

By adopting an approach used to prove Theorem 15 of [1], one can easily prove the following:

THEOREM B. *Let $P(x)$ and $Q(x)$ be any polynomials of degrees at most n . Then there is a constant $c > 0$ for which*

$$\left\| \exp(-e^x) - \frac{P(x)}{Q(x)} \right\|_{L_\infty(0,\infty)} \geq \exp\left(\frac{-cn}{\log \log n}\right).$$

We take this opportunity to correct the following printing errors:

Page	Line	Should read	For
444	13	\leq	\times
449	-4	$\rho (< 1)$ and regular growth	ρ
450	5	$2d_1 d_2 \dots d_n d_{2n+1}$	$2d_{2n+1}$
453	11	$<$	\leq

REFERENCE

1. PAUL ERDÖS, D. J. NEWMAN, and A. R. REDDY, 'Rational approximation (II)', *Advances in Math.*, to appear.

*Hungarian Academy of Sciences
Budapest*

*Institute for Advanced Study
Princeton, N.J. 08540*

RATIONAL APPROXIMATION ON THE POSITIVE REAL AXIS

By PAUL ERDÖS and A. R. REDDY

[Received 31 October 1973—Revised 18 June 1974]

Introduction

Rational Chebyshev approximation to reciprocals of certain entire functions by reciprocals of polynomials on the positive real axis has recently attracted the attention of many mathematicians. By developing certain new methods of approach we successfully attacked ([3]–[6]) some of the related problems. This paper is a continuation of our earlier papers ([3]–[6]). The results of this paper improve and extend some of the earlier results with simplified proofs (cf. Theorem 3). For a reader interested in this topic, this paper may serve as a guide by illustrating some of the techniques (old ones with refinements, as well as new) which we used to solve some of the very interesting and difficult problems of the field (cf. examples 1, 2, 3 of Theorem 5).

Notation and definitions

Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ be any entire function. As usual, let

$$M(r) = \max_{|z|=r} |f(z)|, \quad m(r) = \max_{n \geq 1} |a_n| r^n = |a_\nu| r^\nu,$$

where $\nu = \nu(r)$ is an increasing function of r . $M(r)$, $m(r)$, and $\nu(r)$ are known as the *maximum modulus*, *maximum term*, and the *rank of the maximum term*, respectively. If there exists more than one term which is equal to the maximum term, then we take the one with the largest index. $S_n(z)$ denotes the n th partial sum of $f(z)$. π_n denotes the class of ordinary polynomials of degree at most n , $\pi_{m,n}$ denotes the class of all rational functions of the form $r_{m,n} = p_m/q_n$, where $p_m \in \pi_m$, $q_n \in \pi_n$. Throughout our work we denote ($k \geq 1$):

$$l_k(x) = l_{k-1}[\log x], \quad l_0(x) = x;$$

$$e_k(x) = e_{k-1}[\exp x], \quad e_0(x) = x;$$

$$n(l_1 n)(l_2 n)(l_3 n) \dots (l_{k+1} n)^2 = A(n);$$

$$(l_1 n)(l_2 n)(l_3 n) \dots (l_k n)^{1+\varepsilon} = B(n);$$

$$(l_1 n)(l_2 n)(l_3 n) \dots (l_k n) = D(n).$$

As usual we write

$$(1) \quad \lambda_{0,n}(f^{-1}) \equiv \lambda_{0,n} \equiv \inf_{p \in \pi_n} \left\| \frac{1}{f(x)} - \frac{1}{p(x)} \right\|_{[0,\infty)},$$

where $\|\cdot\|$ is the uniform norm on $[0, \infty)$. As usual we define the order ρ of $f(z)$ as follows ([2], p. 8). The entire function $f(z)$ is of order ρ if

$$\limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log r} = \rho \quad (0 \leq \rho \leq \infty).$$

If ρ is positive and finite, then we define the type τ and the lower type ω , corresponding to the order ρ , as follows:

$$(2) \quad \limsup_{r \rightarrow \infty} r^{-\rho} \log M(r) = \tau, \quad \liminf_{r \rightarrow \infty} r^{-\rho} \log M(r) = \omega$$

$$(0 < \rho < \infty, 0 \leq \omega \leq \tau \leq \infty).$$

It is known ([2], p. 13) that for functions of finite order we can replace $\log M(r)$ by $\log m(r)$ in the above formulae. That is,

$$\limsup_{r \rightarrow \infty} \frac{\log \log m(r)}{\log r} = \rho \quad (0 \leq \rho \leq \infty),$$

$$(2') \quad \limsup_{r \rightarrow \infty} r^{-\rho} \log m(r) = \tau, \quad \liminf_{r \rightarrow \infty} r^{-\rho} \log m(r) = \omega$$

$$(0 < \rho < \infty, 0 \leq \omega \leq \tau \leq \infty).$$

If $f(z)$ is of order zero, then we define as in [11], p. 145, the logarithmic order $\rho_l = \Lambda + 1$, and if Λ is strictly positive and finite then the corresponding logarithmic types are defined as follows:

$$\limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \limsup_{r \rightarrow \infty} \frac{\log \log m(r)}{\log \log r} = \Lambda + 1 \quad (0 \leq \Lambda \leq \infty),$$

$$\limsup_{r \rightarrow \infty} \frac{\log M(r)}{(\log r)^{\Lambda+1}} = \limsup_{r \rightarrow \infty} \frac{\log m(r)}{(\log r)^{\Lambda+1}} = \tau_l,$$

$$\liminf_{r \rightarrow \infty} \frac{\log M(r)}{(\log r)^{\Lambda+1}} = \liminf_{r \rightarrow \infty} \frac{\log m(r)}{(\log r)^{\Lambda+1}} = \omega_l \quad (0 < \Lambda < \infty, 0 \leq \omega_l \leq \tau_l \leq \infty).$$

It is also known ([20], p. 45) that if $f(z) = \sum_{k=0}^{\infty} a_k z^k$ is of order ρ ($0 < \rho < \infty$), type τ , and lower type ω ($0 < \omega \leq \tau < \infty$), then

$$\limsup_{n \rightarrow \infty} (n/\rho e) |a_n|^{\rho/n} = \tau,$$

and

$$(3) \quad \liminf_{p \rightarrow \infty} (n_p/\rho e) |a_{n_p}|^{\rho/n_p} \geq \omega,$$

for a sequence of numbers n_p satisfying the condition

$$(4) \quad \limsup_{p \rightarrow \infty} (n_{p+1}/n_p) \leq x_1/x_2,$$

where x_1 is the greatest and x_2 the smallest root of the equation

$$(5) \quad x \log(x/\epsilon) + (\omega/\tau) = 0.$$

LEMMA 1 ([7], pp. 534-35). Let $p(x)$ be any algebraic polynomial of degree at most n . If this polynomial is bounded by M on an interval of total length l contained in $[-1, 1]$, then in $[-1, 1]$,

$$(6) \quad |p(x)| \leq M |T_n(4l^{-1} - 1)|$$

where $2T_n(x) = (x + \sqrt{(x^2 - 1)})^n + (x - \sqrt{(x^2 - 1)})^n$.

LEMMA 2 ([20], p. 34). Let $f(z) = \sum_0^\infty a_k z^k$ be any entire function of finite order ρ . Then for any $\epsilon > 0$, and all sufficiently large $r \geq r_0(\epsilon)$, we have

$$M(r) \leq m(r)r^{\rho+\epsilon}.$$

LEMMA 3 ([10], Problem [1], part I). Let

$$(7) \quad f(x) = 1 + \sum_{j=1}^{\infty} \frac{x^j}{d_1 d_2 d_3 \dots d_j} \quad (d_{j+1} > d_j > 0, j \geq 1).$$

Then for $x = d_n$, the n th term of the series (7) becomes the maximum term.

For the detailed discussion of our results, we need the following known results.

THEOREM I ([8], Theorem 6). Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ be an entire function of order ρ , type τ , and lower type ω ($0 < \omega \leq \tau < \infty$), with $a_0 > 0$ and $a_k \geq 0$ for all $k \geq 1$. Then

$$\limsup_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} < 1.$$

THEOREM II ([12], Theorem 7'). Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be an entire function satisfying the assumptions that $0 < \Lambda < \infty$ and $0 < \omega_1 \leq \tau_1 < \infty$. Then

$$\limsup_{n \rightarrow \infty} (\lambda_{0,n})^{n^{-1-(\Lambda/\Lambda)}} < 1.$$

THEOREM III ([13], Theorem D). Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be any entire function of order ρ ($0 < \rho < \infty$), type τ , and lower type ω , with the assumption that $\tau < \theta\omega$ for a $\theta < 2$ and $0 < \omega \leq \tau < \infty$. Then

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \geq (\omega/\tau 2^{2\rho+1})^{1/\rho 2^n}.$$

THEOREM IV ([3], Theorem 1). Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be any entire function. Then for each $\epsilon > 0$ there exist infinitely many n such that

$$\lambda_{0,n} \leq \exp(-n(\log n)^{-1-\epsilon}).$$

THEOREM V ([3], Theorem 2). *Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ be any entire function of infinite order with non-negative coefficients. Then for each $\varepsilon > 0$ there exist infinitely many n for which*

$$\lambda_{0,n} \geq \exp(-\varepsilon n).$$

THEOREM VI ([18], Theorem). *Let $f(x) = e^x$. Then*

$$\lim_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} = \frac{1}{3}.$$

Careful observation of the above theorems naturally leads to the following questions.

QUESTION 1. Can one obtain under the assumptions of Theorem I the fact that

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} > 0?$$

QUESTION 2. Is it possible to improve the upper bound and provide a simple proof to Theorem I?

QUESTION 3. What conclusion do we get by dropping the assumptions on the logarithmic types in Theorem II?

QUESTION 4. Is it possible to prove Theorem III without the assumption that $\tau < \theta\omega$?

QUESTION 5. Is it possible to replace $(\log n)^{1+\varepsilon}$ by $(l_1 n)(l_2 n) \dots (l_k n)^{1+\varepsilon}$ for any $k \geq 1$ in Theorem IV?

QUESTION 6. Given an $\varepsilon_n \geq (\log \log n)^{-1}$ can we replace ε in Theorem V by ε_n ?

QUESTION 7. Are there any other functions besides e^x for which we get, for a $\psi(n)$ which tends to infinity,

$$\lim_{n \rightarrow \infty} (\lambda_{0,n})^{1/\psi(n)} = \delta \quad (0 < \delta < 1)?$$

These questions motivated the work of this paper and in it we answer all of them.

New results

THEOREM 1. *Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be any entire function. Then for each $\varepsilon > 0$ and any $k \geq 1$, there exist infinitely many n such that*

$$(8) \quad \lambda_{0,n} \leq \exp(-n/(l_1 n)(l_2 n) \dots (l_k n)^{1+\varepsilon}).$$

Proof. If $f(z) = \sum_{k=0}^{\infty} a_k z^k$ is entire, then $\lim_{n \rightarrow \infty} |a_n|^{1/n} = 0$. Let $u_n = a_n^{-1/n}$. Then $u_n \rightarrow \infty$. Now it is easy to observe from the

convergence of

$$\prod_{j=c_{k+1}(2)}^{\infty} (1 + [A(j)]^{-1}),$$

that there exist arbitrarily large values of n for which for each $l > 0$,

$$(9) \quad u_{n+l} > u_n \prod_{t=1}^l (1 + [A(n+t)]^{-1}).$$

From (9) it follows, with $l = n$, that

$$(10) \quad u_{2n} > u_n (1 + n[2A(n)]^{-1}).$$

Given any $\varepsilon > 0$, we can show now that there exist infinitely many n such that

$$(11) \quad \left\| \frac{1}{f(x)} - \frac{1}{S_{2n}(x)} \right\|_{[0, \infty)} < \exp(-2n[B(2n)]^{-1}).$$

By the definition of $\lambda_{0,n}$ (8) follows from (11). To prove (11), observe that, on the one hand, we have for all $x \geq 0$,

$$0 \leq \frac{1}{S_{2n}(x)} - \frac{1}{f(x)} \leq \frac{1}{S_{2n}(x)} \leq \frac{1}{a_n x^n}.$$

Now for any given $\varepsilon > 0$,† let $x \geq u_n(1 + [B(n)]^{-1})$. Then

$$(12) \quad a_n x^n \geq (1 + [B(n)]^{-1})^n \geq \exp(2n[B(2n)]^{-1}).$$

On the other hand, let $x < u_n(1 + [B(n)]^{-1})$. Then for all $n \geq n_1$,

$$(13) \quad 0 \leq \frac{1}{S_{2n}(x)} - \frac{1}{f(x)} = \frac{f(x) - S_{2n}(x)}{f(x)S_{2n}(x)} \leq a_0^{-2} \sum_{k=2n+1}^{\infty} a_k x^k.$$

By (9) and (10), we have for all $k > 2n$,

$$(14) \quad a_k < u_n^{-k} (1 + n[2A(n)]^{-1})^{-k}.$$

Thus, from (13) and (14), for $x < u_n(1 + [B(n)]^{-1})$, we obtain

$$(15) \quad a_0^{-2} \sum_{k=2n+1}^{\infty} a_k x^k \leq a_0^{-2} \sum_{k=2n+1}^{\infty} \left(\frac{1 + [B(n)]^{-1}}{1 + n[2A(n)]^{-1}} \right)^k.$$

A simple calculation based on (15) gives us

$$(16) \quad a_0^{-2} \sum_{k=2n+1}^{\infty} a_k x^k \leq \exp(-2n[B(2n)]^{-1}).$$

Inequality (11) now follows from (12) and (16).

THEOREM 2. *For every large $c > 0$ and $k \geq 1$ there is an entire function of infinite order with non-negative coefficients for which there exist infinitely*

† ε may not be the same at each occurrence.

many n such that

$$\lambda_{0,n} \geq \exp(-cn/(l_1 n)(l_2 n) \dots (l_k n)).$$

Proof. Let $f(x) = e_{k+1}(x)$ ($k \geq 1$), and let us suppose that for all large n ,

$$(17) \quad \left\| \frac{1}{f} - \frac{1}{p} \right\|_{[0, \infty)} < \exp(-cn[D(n)]^{-1}).$$

Let $x = l_k(n[D(n)]^{-1}) = r_1(k)$, say ($k \geq 1$). At this point

$$f(x) = e_{k+1}(x) = \exp(n[D(n)]^{-1}).$$

Then, by (17),

$$(18) \quad |p| < \exp(2n[D(n)]^{-1}).$$

But at $x = l_k(ne^2[D(n)]^{-1}) = r_2(k)$, say,

$$(19) \quad f(x) = e_{k+1}(x) = \exp(ne^2[D(n)]^{-1}).$$

By applying Lemma 1 to (18) we get for the interval $[0, r_2]$

$$(20) \quad |p| \leq \exp\left(\frac{2n}{(l_1 n)(l_2 n) \dots (l_k n)}\right) T_n \left(\frac{2l_k(ne^2[D(n)]^{-1})}{l_k(n[D(n)]^{-1})} - 1\right) \\ \times \exp(2n[D(n)]^{-1}) \exp(4n[D(n)]^{-1}).$$

From (19) and (20), it is easy to see that

$$\left\| \frac{1}{f(x)} - \frac{1}{p(x)} \right\|_{[0, r_2]} \geq \exp(-cn[D(n)]^{-1}),$$

for some constant c , which contradicts our earlier assumption (17). Hence the required result is proved.

THEOREM 3. Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be any entire function of order ρ ($0 < \rho < \infty$), type τ , and lower type ω ($0 < \omega \leq \tau < \infty$). Then

$$0 < (e\omega^2/e^{2\omega/\tau(e+1)}\tau^2(e+1)4^\rho)^{x_1/x_2} \leq \liminf_{n \rightarrow \infty} (\lambda_{0,n})^{\rho/n} \\ \leq \limsup_{n \rightarrow \infty} (\lambda_{0,n})^{\rho/n} \leq \exp(-\omega/(e+1)\tau) < 1,$$

where x_1 is the greatest and x_2 the smallest root of equation (5).

Proof. If $f(z)$ is an entire function of order ρ ($0 < \rho < \infty$) and type τ ($0 < \tau < \infty$), then, for each $\varepsilon > 0$, there is an $n_2 = n_2(\varepsilon)$ such that, for all $n \geq n_2(\varepsilon)$, we have ([2], p. 11)

$$(21) \quad |a_n| \leq (\rho e \tau (1 + \varepsilon) / n)^{n/\rho},$$

As earlier, we have

$$(22) \quad 0 \leq \frac{1}{S_{n_2}(x)} - \frac{1}{f(x)} \leq a_0^{-2}(f(x) - S_{n_2}(x)) \leq a_0^{-2} \sum_{k=n_2+1}^{\infty} a_k x^k \\ \leq a_0^{-2} \sum_{k=n_2+1}^{\infty} a_k r^k \quad (0 \leq x \leq r).$$

On the other hand, for each $r > 0$, we can find ([2], p. 12) an $n_3 = n_3(r)$ such that for all $n \geq n_3^\dagger$ and $x \geq r$,

$$(23) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq \frac{1}{S_n(x)} \leq \frac{1}{m(r)}.$$

Now two possibilities occur in (23): (i) $n_3 > n_2$, or (ii) $n_3 \leq n_2$. If (i) is true, then in (22) we replace $S_{n_2}(x)$ by $S_{n_3}(x)$, that is,

$$(22') \quad 0 \leq \frac{1}{S_{n_3}(x)} - \frac{1}{f(x)} \leq a_0^{-2} \sum_{k=n_2+1}^{\infty} a_k r^k.$$

If (ii) is true, then in (23) we replace $S_{n_3}(x)$ by $S_{n_2}(x)$, that is, for all $x \geq r$,

$$(23') \quad 0 \leq \frac{1}{S_{n_2}(x)} - \frac{1}{f(x)} \leq \frac{1}{m(r)},$$

where

$$(24) \quad r = (n/\rho\tau(e+1)(1+\varepsilon))^{1/\rho}.$$

In either case we choose $n = \max(n_2, n_3)$.[‡] A simple calculation based on (21) and (22') gives us, for $0 \leq x \leq r$,

$$(25) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq a_0^{-2} \sum_{k=n+1}^{\infty} a_k r^k \\ \leq a_0^{-2} \sum_{k=n+1}^{\infty} \left(\frac{\rho e \tau (1 + \varepsilon)}{\rho \tau (e + 1) (1 + \varepsilon)} \right)^{k/\rho} \\ \leq a_0^{-2} \sum_{k=n+1}^{\infty} \left(\frac{e}{e + 1} \right)^{k/\rho} \\ \leq a_0^{-2} \left(\frac{e}{e + 1} \right)^{(n+1)/\rho} \left(\frac{(e + 1)^{1/\rho}}{(e + 1)^{1/\rho} - e^{1/\rho}} \right).$$

On the other hand, from (2') and (23'), for all $r \geq r_3(\varepsilon)$,

$$(26) \quad m(r) \geq \exp[r^\rho \omega (1 - \varepsilon)] = \exp\left(\frac{\omega n (1 - \varepsilon)}{(e + 1) \rho \tau (1 + \varepsilon)} \right) = G\left(\frac{n(1 - \varepsilon)}{(1 + \varepsilon)}, \rho, \omega, \tau \right),$$

[†] n_3 denotes the rank of the maximum term.

[‡] It is easy to verify that $n_3(r) \leq n(r)$ for the value of r given in (24).

where ε is arbitrary. We easily obtain from (25) and (26) that

$$\limsup_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \leq \exp(-\omega/\rho\tau(e+1)) < 1.$$

Now we shall prove the other inequality of Theorem 3. As the coefficients of $f(x)$ are non-negative, we have from (2), for all large $r \geq r_4(\varepsilon)$,

$$(27) \quad 0 \leq f(x) \leq f(r) = M(r) \leq \exp(r^\rho \tau(1+\varepsilon)) \quad (0 \leq x \leq r, r \geq r_4(\varepsilon)).$$

Now from (27) we have, for

$$r = \{\omega n(1+2\varepsilon)^{-1} \rho^{-1} \tau^{-2} (e+1)^{-1}\}^{1/\rho} = H(n, \rho, \omega, \tau),$$

that

$$\begin{aligned} 0 \leq f(x) &\leq f \left[\left(\frac{\omega n}{\tau^2 \rho (e+1)(1+2\varepsilon)} \right)^{1/\rho} \right] \\ &\leq \exp \left(\frac{n\omega(1+\varepsilon)}{(1+2\varepsilon)\tau\rho(e+1)} \right) \\ &< G(n, \rho, \omega, \tau) = \exp \left(\frac{n\omega}{\tau(e+1)\rho} \right) \leq \frac{1}{\lambda_{0,n}} \end{aligned}$$

for all $n \geq n_4$. Next, we take the rational function $r_{0,n}^* = 1/p_n^*$ ($r_{0,n}^* \in \pi_{0,n}$) which gives the best approximation in the sense of (1); that is, for all $n \geq n_5$,

$$(28) \quad \lambda_{0,n} \equiv \left\| \frac{1}{f(x)} - \frac{1}{p_n^*(x)} \right\|_{[0, \infty)}.$$

A simple manipulation based on (28) gives us

$$\begin{aligned} (29) \quad -f^2(x) / \left(f(x) + \frac{1}{\lambda_{0,n}} \right) &\leq p_n^* - f(x) \\ &\leq f^2(x) / \left(\frac{1}{\lambda_{0,n}} - f(x) \right) \quad (0 \leq x \leq H(n, \rho, \omega, \tau)). \end{aligned}$$

Clearly the right-hand side of inequality (29) is monotonic increasing with x . Hence we write

$$(30) \quad \|p_n^* - f(x)\| \leq G(2n, \rho, \omega, \tau) / \left(\frac{1}{\lambda_{0,n}} - G(n, \rho, \omega, \tau) \right) \quad (0 \leq x \leq H(n, \rho, \omega, \tau)).$$

Next, let

$$(31) \quad E_n(f) \equiv \inf_{p_n \in \pi_n} \|p_n(x) - f(x)\|_{[0, H(n, \rho, \omega, \tau)]}.$$

From (30) and (31) we get, for all $n \geq n_6$,

$$(32) \quad E_n(f) \leq G(2n, \rho, \omega, \tau) / \left(\frac{1}{\lambda_{0,n}} - G(n, \rho, \omega, \tau) \right).$$

To obtain a lower bound for E_n we use a result of Bernstein ([1], p. 10) which gives us, in the interval $[0, H(n, \rho, \omega, \tau)]$,

$$(33) \quad E_n \geq \left(\frac{n\omega}{\tau^2(e+1)\rho(1+2\varepsilon)} \right)^{(n+1)/\rho} \frac{a_{n+1}}{2^{2n+1}}.$$

From (32) and (33) we get

$$(34) \quad \frac{a_{n+1}[H(n, \rho, \omega, \tau)]^{n+1}}{2^{2n+1}} \leq G(2n, \rho, \omega, \tau) / \left(\frac{1}{\lambda_{0,n}} - G(n, \rho, \omega, \tau) \right)$$

for all $n \geq n_7$.

From (3) we have, for a sequence of values of $n = n_p + 1$ satisfying the assumption (4),

$$(35) \quad a_{n_p+1} \geq (\rho e \omega (1 - \varepsilon) / (n_p + 1))^{(n_p+1)/\rho}.$$

From (34) and (35) we get

$$(36) \quad \left(\frac{\rho e \omega (1 - \varepsilon)}{n_p + 1} \right)^{(n_p+1)/\rho} \frac{[H(n_p, \rho, \omega, \tau)]^{n_p+1}}{2^{2n_p+1}} \leq G(2n, \rho, \omega, \tau) / \left(\frac{1}{\lambda_{0,n}} - G(n, \rho, \omega, \tau) \right).$$

It is easy to obtain from (36) that

$$(37) \quad \frac{1}{\lambda_{0,n_p}} \leq G(n_p, \rho, \omega, \tau) + \frac{G(2n_p, \rho, \omega, \tau)(n_p + 1)^{(n_p+1)/\rho} 2^{2n_p+1}}{[\{\rho e \omega (1 - \varepsilon)\}^{1/\rho} H(n_p, \rho, \omega, \tau)]^{n_p+1}} \\ \leq 2 \left[\exp\left(\frac{2n_p \omega}{\tau(e+1)\rho}\right) \right] \left(\frac{4^\rho \tau^2 \rho (e+1)(1+2\varepsilon)}{\omega^2(1-\varepsilon)e} \right)^{(n_p+1)/\rho} \left(\frac{n_p + 1}{n_p} \right)^{(n_p+1)/\rho}.$$

Now by adopting the technique used on p. 373 of [19] we can easily obtain from (37) and (4) the fact that

$$(38) \quad \liminf_{n \rightarrow \infty} (\lambda_{0,n})^{\rho/n} \geq \left(\frac{e\omega^3}{e^{2\omega/\tau(e+1)} \tau^2(e+1)4^\rho} \right)^{\pi_1/\alpha_2}.$$

REMARKS. (1) Under the assumptions of Theorem 3, Reddy ([16]) has recently obtained the following sharper result

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \geq (2^{2+1/\rho} \tau^{1/\rho} \omega^{-1/\rho} - 1)^{-2}.$$

(2) There exist entire functions which fail to satisfy the assumptions of Theorem 3, but for which we can still find two constants c_1 and c_2 ($0 < c_1 \leq c_2 < 1$) such that

$$0 < c_1 \leq \liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \leq \limsup_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \leq c_2 < 1.$$

EXAMPLE 1. Let

$$f(x) = 1 + \sum_{k=2}^{\infty} \left(\frac{\log n}{n} \right)^n x^n.$$

This is an entire function of order $\rho = 1$ and type $\tau = \infty$. For this function the maximum module $M(r)$ is given by

$$(39) \quad f(r) = M(r) \sim \exp\left(\frac{r}{e} \log \frac{r}{e}\right).$$

This function clearly satisfies the growth condition (3.1) of [8]; hence there exists a constant $q > 1$ for which

$$(40) \quad \limsup_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} = \frac{1}{q} < 1.$$

From (4) it is easy to see that, for all large $n \geq n_8(\varepsilon)$,

$$(41) \quad q_1^n = [q(1-\varepsilon)]^n \leq 1/\lambda_{0,n}.$$

From (39) we have

$$0 \leq f(x) \leq f(r) < \exp(r \log(r/e)).$$

Let

$$(42) \quad r \log(r/e) = \frac{1}{2} n \log q_1.$$

That is,

$$r \sim \frac{n \log q_1}{2 \log(\frac{1}{2} n \log q_1)}.$$

From (41) and (42) we get, for all $n \geq n_9$,

$$0 \leq f(x) \leq f(r) \leq \exp(r \log(r/e)) \leq q_1^{n/2} < 1/\lambda_{0,n}.$$

Now proceeding exactly as in the proof of the second part of Theorem 3 we get, for the value of r given in (42) and for all $n \geq n_{10}$,

$$(43) \quad \left(\frac{n}{\log(\frac{1}{2} n \log q_1)} \right)^{n+1} \frac{a_{n+1}}{2^{2n+1}} \leq q_1^n / \left(\frac{1}{\lambda_{0,n}} - q_1^{n/2} \right).$$

From (43) it is easy to see that

$$\begin{aligned} \frac{1}{\lambda_{0,n}} &\leq \frac{2^{2n+1} q_1^n [\log(\frac{1}{2} n \log q_1)]^{n+1}}{n^{n+1} a_{n+1}} + q_1^{n/2} \\ &\leq 2^{2n+1} q_1^n \left(\frac{\log(\frac{1}{2} n \log q_1)}{n \log(n+1)} \right)^{n+1} (n+1)^{n+1} + q_1^{n/2}. \end{aligned}$$

In other words,

$$(44) \quad \frac{1}{\lambda_{0,n}} \leq 2^{2n+2} q_1^n \left(\frac{n+1}{n} \right)^{n+1} \left(\frac{\log(\frac{1}{2} n) + \log \log q_1}{\log(n+1)} \right)^{n+1}.$$

A simple manipulation based on (44) gives us, if we take q_1 to be very close to q ,

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \geq 1/4q.$$

EXAMPLE 2. Let

$$f(x) = 1 + \sum_{n=2}^{\infty} \left(\frac{x}{n \log n} \right)^n.$$

This is an entire function of order $\rho = 1$ and type $\tau = 0$. For this function

$$(45) \quad f(r) \sim \exp\left(\frac{r}{e \log(r/e)}\right).$$

$f(r)$ satisfies the growth condition (3.1) given in [8]; hence there exists a $q > 1$ such that

$$(46) \quad \limsup_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} = \frac{1}{q} < 1.$$

As before from (46) we can find, for all $n \geq n_{11}(\varepsilon)$, a $q_1 < q$ such that

$$(47) \quad \lambda_{0,n} \leq q_1^{-n}.$$

Let

$$(48) \quad \frac{r}{2 \log(r/e)} = \frac{1}{2} n \log q_1.$$

Then for some $c > 0$, $r \sim n \log(cn \log q_1)$. From (45), (47), and (48) we obtain, for all $n \geq n_{12}$,

$$0 \leq f(x) \leq f(r) \leq \exp\left(\frac{(1+\varepsilon)r}{e \log(r/e)}\right) \leq q_1^{n/2} < \frac{1}{\lambda_{0,n}}$$

and

$$(49) \quad \frac{n^{(n+1)} [\log(cn \log q_1)]^{n+1} a_{n+1}}{2^{2n+1}} \leq q_1^n / \left(\frac{1}{\lambda_{0,n}} - q_1^{n/2} \right).$$

From (49) we get as before, for $a_n = (n \log n)^{-n}$,

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{1/n} \geq 1/4q.$$

THEOREM 4. Let $f(z) = 1 + \sum_{k=1}^{\infty} a_k z^k$, $a_k = (d_1 d_2 \dots d_k)^{-1}$ with $d_{k+1} > d_k > 0$ ($k \geq 1$), be an entire function of finite order ρ . Then for any $\varepsilon > 0$ and all large $n \geq n_{13}(\varepsilon)$, we have

$$(50) \quad \frac{d_1 d_2 d_3 \dots d_n}{2^{4n} d_n^{2(\rho+\varepsilon)} d_{n+1} d_{n+2} \dots d_{2n}} \leq \lambda_{0,2n-1} \leq \frac{d_1 d_2 d_3 \dots d_n d_{2n+1}}{d_{n+1} d_{n+2} \dots d_{2n} (d_{2n+1} - d_{2n})}.$$

Proof. The second half of (50) follows from the proof of Theorem 5 of [4].

To prove the first half of (50) observe that, for $0 \leq x \leq r = d_n$, Lemmas 2 and 3 hold and that, for all $n \geq \hat{n}(\varepsilon)$,

$$(51) \quad 0 \leq f(x) \leq M(r) \leq m(r)r^{\rho+\varepsilon} \\ = m(d_n)d_n^{\rho+\varepsilon} \\ = \frac{d_n^n d_n^{\rho+\varepsilon}}{d_1 d_2 \dots d_n} < \frac{d_{n+1} d_{n+2} \dots d_{2n} (d_{2n+1} - d_{2n})}{2 d_{2n+1}} = \frac{(d_{2n+1} - d_{2n})}{2 J(d_n) d_{2n+1}}.$$

This follows because when $x = d_n$ we know from Lemma 3 that the n th term becomes the maximum term, that is,

$$m(d_n) = d_n^n / d_1 d_2 \dots d_n.$$

As before we choose $p^*(x) \in \pi_{2n-1}$ such that $p^*(x)$ gives the best approximation to $f(x)$ in the sense of (1), that is,

$$(52) \quad \lambda_{0,2n-1} \geq \left\| \frac{1}{f(x)} - \frac{1}{p^*(x)} \right\|_{[0,\infty)} \quad (p^* \in \pi_{2n-1}).$$

A simple calculation based on (52) gives us (as in the proof of Theorem 3)

$$(53) \quad \|f - p_{2n-1}^*\| < \{f(x)\}^2 / \left(\frac{1}{\lambda_{0,2n-1}} - f(x) \right) \quad (0 \leq x \leq r = d_n).$$

From (51) and (53) we obtain

$$(54) \quad \|f - p_{2n-1}^*\| \\ \leq d_n^{2n} d_n^{2(\rho+\varepsilon)} / (d_1 d_2 \dots d_n)^2 \left(\frac{1}{\lambda_{0,2n-1}} - \frac{d_n^n d_n^{\rho+\varepsilon}}{d_1 d_2 \dots d_n} \right) \quad (0 \leq x \leq d_n).$$

Let

$$(55) \quad E_{2n-1}(f) = \min_{g_{2n-1} \in \pi_{2n-1}} \|f - g_{2n-1}\|_{[0,d_n]}.$$

From (54) and (55) we get

$$(56) \quad E_{2n-1}(f) \\ \leq d_n^{2n} d_n^{2(\rho+\varepsilon)} / (d_1 d_2 \dots d_n)^2 \left(\frac{1}{\lambda_{0,2n-1}} - \frac{d_n^n d_n^{\rho+\varepsilon}}{d_1 d_2 \dots d_n} \right) \quad (0 \leq x \leq d_n).$$

To obtain a lower bound for $E_{2n-1}(f)$ we use (as before) a result of Bernstein ([1], p. 10), which gives us for $x = d_n$,

$$(57) \quad E_{2n-1}(f) \geq a_{2n} d_n^{2n} / 2^{2n} 2^{2n-1}.$$

From (56) and (57) we get, for all $n \geq n_{14}$,

$$(58) \quad J(d_n) \leq 2^{2n-1} d_n^{2(\rho+\varepsilon)} / \left(\frac{1}{\lambda_{0,2n-1}} - \frac{d_n^n d_n^{\rho+\varepsilon}}{d_1 d_2 \dots d_n} \right).$$

From (58) it is easy to calculate that

$$\frac{1}{\lambda_{0,2n-1}} \leq \frac{2^{4n} d_n^{2(\rho+\varepsilon)}}{J(d_n)}.$$

Therefore for all large n , we have

$$\frac{d_1 d_2 d_3 \dots d_n}{2^{4n} d_n^{2(\rho+\varepsilon)} d_{n+1} d_{n+2} \dots d_{2n}} \leq \lambda_{0,2n-1} \leq \frac{d_1 d_2 \dots d_n d_{2n+1}}{d_{n+1} d_{n+2} \dots d_{2n} (d_{2n+1} - d_{2n})}.$$

EXAMPLES. (1) Let

$$f(x) = 1 + \sum_{n=1}^{\infty} \frac{x^n}{2^{\log 2} 3^{\log 3} \dots n^{\log n}}.$$

For this function we get from (50)

$$\lim_{n \rightarrow \infty} (\lambda_{0,n})^{1/(n \log n)} = \frac{1}{2}.$$

It is interesting to note that this function fails to satisfy the assumptions of Theorem 7 of [8] and Theorem 7' of [12], because $\rho_1 = \Lambda + 1 = \infty$. But our present method which is much simpler than the methods used in [8] and [12] gives us more precise information.

(2) Let

$$f(x) = 1 + \sum_{n=1}^{\infty} \frac{x^n}{2^2 3^3 4^4 \dots n^n}.$$

This function also fails to satisfy the assumptions of Theorem 7 of [8] and Theorem 7' of [12], because in this case $\Lambda = 1$, that is, $\rho_1 = 2$ and $\tau_1 = 0$. For this function we get by (50)

$$\lim_{n \rightarrow \infty} (\lambda_{0,n})^{1/n^2 \log n} = e^{-1/4}.$$

(3) Let

$$f(x) = 1 + \sum_{n=1}^{\infty} \frac{x^n}{\delta^{2^n}} \quad (1 < \delta < \infty).$$

For this function $\Lambda = 0$, whence it also fails to satisfy the assumptions of Theorem 7 of [8] and Theorem 7' of [12]; but we obtain by (50)

$$\lim_{n \rightarrow \infty} (\lambda_{0,n})^{2^{-n-1}} = 1/\delta.$$

THEOREM 5. Let $f(x) = 1 + \sum_{k=1}^{\infty} (d_1 d_2 \dots d_k)^{-1} x^k$ ($d_{k+1} > d_k > 0$, $k \geq 1$) be an entire function of infinite order. Then for each $\varepsilon > 0$, there exist infinitely many n for which

$$\lambda_{0,2n-1} \geq [K(d_n)]^{1+\varepsilon}.$$

Proof. As before when $x = d_n$, $d_n^n/d_1 d_2 d_3 \dots d_n$ becomes the maximum term of f . Let $t_n = x^n/d_1 d_2 \dots d_n$. Then

$$(59) \quad f(x) = t_n + t_{n-1} \left(1 + \frac{t_{n-2}}{t_{n-1}} + \frac{t_{n-3}}{t_{n-2}} + \dots + t_0 \right) \\ + t_{n+1} \left(1 + \frac{t_{n+2}}{t_{n+1}} + \frac{t_{n+3}}{t_{n+2}} + \dots \right).$$

But

$$\frac{t_{n+2}}{t_{n+1}} = \frac{x}{d_{n+2}} < \frac{d_{n+1}}{d_{n+2}}, \\ \frac{t_{n+3}}{t_{n+2}} = \frac{x^2}{d_{n+2} d_{n+3}} < \left(\frac{d_{n+1}}{d_{n+2}} \right)^2,$$

and so on. Hence

$$(60) \quad \left| \frac{t_{n+2}}{t_{n+1}} + \frac{t_{n+3}}{t_{n+2}} + \dots \right| < \frac{d_{n+1}}{d_{n+2} - d_{n+1}}.$$

Similarly

$$\frac{t_{n-2}}{t_{n-1}} \leq \left(\frac{d_{n-1}}{d_n} \right), \quad \frac{t_{n-3}}{t_{n-2}} \leq \left(\frac{d_{n-1}}{d_n} \right)^2,$$

and so on. Therefore

$$(61) \quad \left| \frac{t_{n-2}}{t_{n-1}} + \frac{t_{n-3}}{t_{n-2}} + \dots \right| < \frac{d_{n-1}}{d_n - d_{n-1}}.$$

From (59), (60), and (61), we obtain

$$(62) \quad f(x) = t_n + t_{n-1}(1 + \varphi) + t_{n+1}(1 + \varphi_1),$$

where

$$|\varphi| < \frac{d_{n-1}}{d_n - d_{n-1}}, \quad |\varphi_1| < \frac{d_{n+1}}{d_{n+2} - d_{n+1}}.$$

From (62) we get, for $x = d_n$,

$$f(x) \leq \frac{d_n^n}{d_1 d_2 d_3 \dots d_n} + t_{n-1}(1 + \varphi) + t_{n+1}(1 + \varphi_1) = [K(d_n)]^{-1/\delta}.$$

For all sufficiently large n we can find an $\varepsilon > 0$ such that

$$(63) \quad f(d_n) \leq [K(d_n)]^{-(1+\varepsilon)/\delta}.$$

Now let us suppose that for all large n and all x ($0 \leq x < \infty$)

$$(64) \quad \left| \frac{1}{f(x)} - \frac{1}{p_{2n-1}(x)} \right| \leq [K(d_n)]^{1+\varepsilon}.$$

From (63) and (64) we get

$$|p| < [K(d_n)]^{-(1+\varepsilon)/4}.$$

Because of the assumption that $f(x)$ is of infinite order for every large $r > 0$ we can find sufficiently large n such that

$$(65) \quad f[d_n(1+r^{-1})] \geq [f(d_n)]^{18}.$$

Therefore from (62) and (65) we get

$$(66) \quad f[d_n(1+r^{-1})] \geq [K(d_n)]^{-2(1+\varepsilon)}.$$

On the other hand by applying Lemma 1 to $p(x)$ we get

$$(67) \quad p[d_n(1+r^{-1})] \leq [K(d_n)]^{-(1+\varepsilon)/2}.$$

From (66) and (67) we get

$$(68) \quad [K(d_n)]^{1+\varepsilon} \leq [K(d_n)]^{(1+\varepsilon)/2} \{1 - [K(d_n)]^{3(1+\varepsilon)/2}\} \\ \leq \frac{1}{p_n(d_n(1+r^{-1}))} - \frac{1}{f(d_n(1+r^{-1}))},$$

that is,

$$\{K(d_n)\}^{(1+\varepsilon)/2} + \{K(d_n)\}^{3(1+\varepsilon)/2} < 1,$$

which gives the required result, because the conclusion (68) contradicts our earlier assumption (64).

THEOREM 6. Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_0 > 0$, $a_k \geq 0$, $k \geq 1$) be any entire function satisfying the assumption that

$$1 \leq \limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \Lambda + 1 < \infty.$$

Then for any $\varepsilon > 0$,

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{n^{-1-(\Lambda+\varepsilon)}} < 1.$$

Proof. We get from Theorems 1 and 3 of [II]

$$\limsup_{n \rightarrow \infty} \frac{\log n}{\log\{n^{-1} \log a_n^{-1}\}} = \Lambda.$$

From this we easily obtain that, for any $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} |a_n|^{1/n} \exp(n^{1/(\Lambda+\varepsilon)}) = 0.$$

As earlier, let $u_n = a_n^{-1/n}$, and let $h = 1/(\Lambda + \varepsilon)$. Then $u_n \exp(-n^h) \rightarrow \infty$. This implies that there exist infinitely many n for which

$$(69) \quad \frac{u_{n+l}}{\exp[(n+l)^h]} \geq \frac{u_n}{\exp(n^h)} \quad (l = 0, 1, 2, \dots).$$

Now let

$$x \geq \theta^{n^h} u_n \quad (1 < \theta < \exp(2^{h-1})).$$

Then

$$(70) \quad S_{2n}(x) \geq a_n x^n \geq a_n u_n^n \theta^{n^{h+1}} = \theta^{n^{h+1}}.$$

Hence, as usual we get from (70), for $x \geq \theta^{n^h} u_n$,

$$(71) \quad 0 \leq \frac{1}{S_{2n}(x)} - \frac{1}{f(x)} \leq \frac{1}{S_{2n}(x)} \leq \frac{1}{a_n x^n} \leq \theta^{-n^{h+1}}.$$

On the other hand, let

$$x < u_n (\theta)^{n^h}.$$

Then as before it is easy to note from (69) that, for any $k > 2n$,

$$|a_k| < u_n^{-k} \exp(k(n^h - k^h)).$$

This formula implies that

$$\begin{aligned} \sum_{k=2n+1}^{\infty} a_k x^k &\leq \sum_{k=2n+1}^{\infty} \{\exp k(n^h - k^h)\} \theta^{n^h k} \\ &\leq \sum_{k=2n+1}^{\infty} \left(\frac{(\theta e)^{n^h}}{\exp k^h} \right)^k \\ &\leq \left(\frac{(\theta e)^{n^h}}{\exp(2n)^h} \right)^{2n+1} \left(\frac{\exp(2n)^h}{\exp(2n)^h - (\theta e)^{n^h}} \right). \end{aligned}$$

From this we have

$$\begin{aligned} (72) \quad \lambda_{0,2n} &\leq \left\| \frac{1}{f(x)} - \frac{1}{p_{2n}(x)} \right\| \\ &\leq a_0^{-2} \sum_{k=2n+1}^{\infty} a_k x^k \\ &\leq \left(\frac{(\theta e)^{n^h}}{\exp(2n)^h} \right)^{2n+1} a_0^{-2} \left(\frac{\exp(2n)^h}{\exp(2n)^h - (\theta e)^{n^h}} \right). \end{aligned}$$

From (71) and (72), with $\theta e < \exp(2^h)$, we easily obtain that

$$\liminf_{n \rightarrow \infty} (\lambda_{0,2n})^{1/(2n)^{h+1}} < e^{-1} (\theta e)^{1/(2^h)} < 1.$$

THEOREM 7. Let $f(z) = a_0 + \sum_{k=1}^{\infty} a_{n_k} z^{n_k}$, with $a_0 > 0$, $a_{n_k} > 0$ ($k \geq 1$), and $\liminf_{k \rightarrow \infty} (n_{k+1}/n_k) \geq \beta > 1$, be an entire function of finite order ρ . Then for any α ($0 < \alpha < 1$) and any $\varepsilon > 0$, we have

$$\liminf_{n \rightarrow \infty} (\lambda_{0,n})^{(1-\alpha)(\rho+\varepsilon)/n} \leq \beta^{-1}.$$

The proof of this result is very similar to the proof of the preceding theorem. Hence the details are left to the reader.

Concluding remarks

Theorem 3 of this paper answers questions 1, 2, and 4 in the affirmative. The answer to question 5 follows from Theorem 1. Question 3 is resolved in Theorem 6. The examples given at the end of Theorem 4 answer question 7. Question 6 must be answered in the negative; this follows from Theorem 4.

It may be of some interest to know whether it is possible to obtain a lower bound for

$$\limsup_{n \rightarrow \infty} (\lambda_{0,n})^{n^{-1-n(\Lambda)}} \quad (\text{cf. Theorem II}).$$

This has been solved in [15]. We have proved in [15], under the assumptions of Theorem II, that

$$\limsup_{n \rightarrow \infty} (\lambda_{0,n})^{n^{-1-n(\Lambda)}} \geq \exp\left(\frac{-\Lambda}{(\Lambda+1)} \left(\frac{1}{(\Lambda+1)\tau}\right)^{1/\Lambda}\right).$$

It is natural to ask whether we can do much better by using the general rational functions of the form $p_n(x)/Q_n(x)$ than by using $1/Q_n(x)$ in the above results. We are not able to settle this question in general (see, however, [9] and [17]). But we are able to prove the following theorem.

THEOREM (cf. [14]). *Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$ ($a_k \geq 0$, $k \geq 0$) be any entire function of order ρ ($0 < \rho < \infty$), type τ , and lower type ω ($0 < \omega \leq \tau < \infty$). Then one cannot find algebraic polynomials $p(x)$ and $Q(x)$ with non-negative coefficients and of degree at most n for which*

$$\liminf_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{P(x)}{Q(x)} \right\|_{[0, \infty)} \right\}^{\rho\omega/n\tau} \leq (2\sqrt{2})^{-1}.$$

The examples 1 and 2 of Theorem 3 fail to satisfy the assumptions of the above theorem; but the conclusion of the theorem still holds for these examples, in a slightly different form.

REFERENCES

1. S. N. BERNSTEIN, *Leçons sur les propriétés extrémales et la meilleure approximation des fonctions analytiques d'une variable réelle* (Gauthier-Villars, Paris, 1926).
2. R. P. BOAS, *Entire functions* (Academic Press, New York, 1954).
3. P. ERDÖS and A. R. REDDY, 'Rational approximation to certain entire functions in $[0, +\infty)$ ', *Bull. Amer. Math. Soc.* 79 (1973) 992-93.
4. ———, 'Chebyshev rational approximation to entire functions in $[0, \infty)$ ', *C.R. Acad. Bulgare Sci.* Iliev 60th anniversary volume.
5. ———, 'A note on rational approximation', *Period. Math. Hungar.*, to appear.
6. ———, 'Problems and results in rational approximation', *ibid.*, to appear.
7. ——— and P. TURÁN, 'On interpolation III, interpolatory theory of polynomials', *Ann. of Math.* 41 (1940) 510-53.

8. G. MEINARDUS, A. R. REDDY, G. D. TAYLOR, and R. S. VARGA, 'Converse theorems and extensions in Chebyshev rational approximation to certain entire functions in $[0, +\infty)$ ', *Bull. Amer. Math. Soc.* 77 (1971) 460-61, *Trans. Amer. Math. Soc.* 170 (1972) 171-85.
9. D. J. NEWMAN, 'Rational approximation to e^{-x} ', *J. Approximation Theory* 10 (1974) 301-2.
10. G. PÓLYA and G. SZEGÖ, *Problems and theorems in analysis*, Vol. 1 (Springer-Verlag, Berlin, 1972).
11. A. R. REDDY, 'On entire Dirichlet series of zero order', *Tôhoku Math. J.* 18 (1966) 144-55.
12. — 'Addendum to "Converse theorems and extensions in Chebyshev rational approximation to certain entire functions in $[0, +\infty)$ "', *Trans. Amer. Math. Soc.* 186 (1973) 499-502.
13. — 'A contribution to rational Chebyshev approximation to certain entire functions in $[0, +\infty)$ ', *J. Approximation Theory* 11 (1974) 85-96.
14. — 'A note on rational Chebyshev approximation on the positive real axis', *ibid.* 12 (1974) 201-2.
15. — 'Rational Chebyshev approximation to certain entire functions of zero order on the positive real axis', *ibid.* 15 (1975), to appear.
16. — 'A note on rational approximation on $[0, \infty)$ ', *ibid.* 13 (1974) 489-90.
17. — and O. SHISHA, 'On a class of rational approximations to certain entire functions on the positive real axis—a survey', *ibid.* 12 (1974) 425-34.
18. A. SCHOENHAGE, 'Zur rationalen Approximierbarkeit von e^{-x} über $[0, \infty)$ ', *ibid.* 7 (1973) 395-98.
19. A. F. TIMAN, *Theory of approximation of functions of a real variable*, International series of monographs in pure and applied mathematics 34 (MacMillan, New York, 1963).
20. G. VALIRON, *Lectures on the general theory of integral functions* (Chelsea, New York, 1949).

Hungarian Academy of Sciences
Budapest

Department of Mathematics
Michigan State University
East Lansing, MI 48824