

## Disjoint Edges in Geometric Graphs

N. Alon<sup>1\*</sup> and P. Erdős<sup>2</sup>

<sup>1</sup> Department of Mathematics, Sackler Faculty of Exact Sciences,  
Tel Aviv University, Ramat Aviv, Tel Aviv, Israel

<sup>2</sup> Mathematical Institute of the Hungarian Academy of Sciences,  
Budapest, Hungary

**Abstract.** Answering an old question in combinatorial geometry, we show that any configuration consisting of a set  $V$  of  $n$  points in general position in the plane and a set of  $6n - 5$  closed straight line segments whose endpoints lie in  $V$ , contains three pairwise disjoint line segments.

A *geometric graph* is a pair  $G = (V, E)$ , where  $V$  is a set of points (= *vertices*) in general position in the plane, i.e., no three on a line, and  $E$  is a set of distinct, closed, straight line segments, called *edges*, whose endpoints lie in  $V$ . An old theorem of the second author [Er] (see also [Ku] for another proof), states that any geometric graph with  $n$  points and  $n + 1$  edges contains two disjoint edges, and this is best possible for every  $n \geq 3$ . For  $k \geq 2$ , let  $f(k, n)$  denote the maximum number of edges of a geometric graph on  $n$  vertices that contains no  $k$  pairwise disjoint edges. Thus, the result stated above is simply the fact  $f(2, n) = n$  for all  $n \geq 3$ . Kupitz [Ku] and Perles [Pe] (see also [AA]) raised the problem of determining or estimating  $f(k, n)$  for  $k \geq 3$ . In particular, they asked if  $f(3, n) \leq O(n)$ . This specific problem, of determining or estimating  $f(3, n)$ , was already mentioned in 1966 by Avital and Hanani [AH], and it seems it was a known problem even before that. In this note we answer this question by proving the following.

**Theorem 1.** *For every  $n \geq 1$ ,  $f(3, n) < 6n - 5$ , i.e., any geometric graph with  $n$  vertices and  $6n - 5$  edges contains three pairwise disjoint edges.*

\* Research supported in part by an Allon Fellowship and by a Bat Sheva de-Rothschild grant.

Before proving this theorem we note that clearly

$$f(3, n) = \binom{n}{2} \quad \text{for } n \leq 5$$

and the best-known lower bound for  $n \geq 6$ , given by Perles [Pe], is

$$f(3, n) \geq \begin{cases} \frac{5}{2}n - \frac{5}{2} & \text{for odd } n \geq 5, \\ \frac{5}{2}n - 4 & \text{for even } n \geq 2. \end{cases} \quad (1)$$

To prove inequality (1) for odd  $n$  consider the geometric graph  $G_n$  whose  $n$  vertices are the  $n-1$  points  $v_j = (\cos(2\pi j/(n-1)), \sin(2\pi j/(n-1)))$ ,  $0 \leq j < n-1$ , together with the additional point  $u = (\varepsilon, \delta)$  where  $\varepsilon$  and  $\delta$  are small numbers chosen so that  $\{v_0, \dots, v_{n-1}, u\}$  is in general position. The edges of  $G_n$  are the  $\frac{5}{2}(n-1)$  line segments

$$\begin{aligned} & \{\{u, v_j\}: 0 \leq j < n-1\} \\ & \cup \{\{v_j, v_{j+(n-3)/2}\}, \{v_j, v_{j+(n-1)/2}\}, \{v_j, v_{j+(n+1)/2}\}: 0 \leq j < n-1\}, \end{aligned}$$

where all indices are reduced modulo  $n-1$ . We can easily check that if  $\varepsilon$  and  $\delta$  are sufficiently small then  $G_n$  contains no three pairwise disjoint edges. Thus  $f(3, n) \geq \frac{5}{2}n - \frac{5}{2}$  for every odd  $n \geq 5$ . For even  $n$ , let  $G_n$  be the geometric graph obtained from  $G_{n+1}$  by deleting one of its vertices of degree 4. Then  $G_n$  has  $\frac{5}{2}n - 4$  edges and contains no three pairwise disjoint edges. This establishes (1). On the other hand, Perles [Pe] showed that every geometric graph whose  $n$  vertices are the vertices of a convex  $n$ -gon in the plane, with more than  $(k-1)n$  edges, contains  $k$  pairwise disjoint edges. In particular, in the convex case  $2n+1$  edges guarantee three pairwise disjoint edges. Comparing this with (1) we conclude that the convex case differs from the general one.

Our final remark before the proof of Theorem 1 is that a special case of one of the results in [AA] implies that, for every  $k = o(\log n)$ ,  $f(k, n) = o(n^2)$ . It is very likely that, for every fixed  $k$ ,  $f(k, n) = O(n)$ , and that, for every  $k = o(n)$ ,  $f(k, n) = o(n^2)$ , but this remains open.

*Proof of Theorem 1.* Let  $G$  be a geometric graph with  $n$  vertices and  $6n-5$  edges. We must show that  $G$  contains three pairwise disjoint edges. It is first convenient to apply an affine transformation on the plane, in order to make all the edges of  $G$  almost parallel to the  $x$ -axis. This is done by first choosing the  $x$ -axis so that any two distinct points of  $G$  have different  $x$ -coordinates, and then, by rescaling the  $y$ -coordinates so that the difference between the  $x$ -coordinates of any two distinct points of  $G$  is at least 1000 times bigger than the difference between their  $y$ -coordinates. Since any affine transformation maps disjoint segments into disjoint segments we may apply the above transformations, and hence may assume that  $G$  satisfies the following:

The small angle between any edge of  $G$  and the  $x$ -axis is less than  $\pi/200$ . (2)

We now define the clockwise derivative and the counterclockwise derivative of an arbitrary geometric graph. Let  $H = (V, E)$  be a geometric graph and let  $e = [u, v]$  be an edge of  $H$ . We say that  $e$  is *clockwise good at  $u$*  if there is another edge  $e' = [u, v']$  of  $H$  such that the directed line  $\overrightarrow{uv'}$  is obtained from  $\overrightarrow{uv}$  by rotating it clockwise around  $u$  by an angle smaller than  $\pi/100$ . If  $e$  is not clockwise good at  $u$ , we say that it is *clockwise bad at  $u$* . The edge  $e = [u, v]$  is *clockwise good* if it is clockwise good at both  $u$  and  $v$ . The *clockwise derivative* of  $H$ , denoted by  $\partial H$ , is the geometric graph whose set of vertices is the set of all vertices of  $H$ , and whose set of edges consists of all clockwise good edges of  $H$ . The notions of an edge  $e = [u, v]$  which is *counterclockwise good at  $u$*  and that of an edge which is *counterclockwise good* are defined analogously. The *counterclockwise derivative* of  $H$ , denoted by  $H\partial$ , is also defined in an analogous manner.

**Claim 1.** *Let  $G = (V, E)$  be a geometric graph with  $n \geq 2$  vertices and  $m$  edges satisfying (2). Then the number of edges of  $\partial G$  is at least  $m - (2n - 2)$ . Similarly, the number of edges of  $G\partial$  is at least  $m - (2n - 2)$ .*

*Proof.* We prove the assertion for  $\partial G$ . The proof for  $G\partial$  is analogous. Let  $v \in V$  be an arbitrary vertex of  $G$ . We claim that the number of edges of the form  $[v, u]$  of  $G$  which are clockwise bad at  $v$  does not exceed 2. Indeed, assume this is false and let  $[v, u_1], [v, u_2], [v, u_3]$  be three such edges. Without loss of generality, assume that the  $x$ -coordinates of  $u_1$  and  $u_2$  lie in the same side of the  $x$ -coordinate of  $v$ . By (2), the angle between  $[v, u_1]$  and  $[v, u_2]$  is smaller than  $\pi/100$ , and hence at least one of these two edges is clockwise good at  $v$ . This contradiction shows that indeed at most two edges of the form  $[v, u]$  are clockwise bad at  $v$ . The same argument shows that if  $u$  is a vertex of  $G$  whose  $x$ -coordinate is maximum or minimum, then there is at most one edge incident with  $u$  which is clockwise bad at  $u$ . Altogether, the total number of clockwise bad edges is bounded by  $2 + 2 \cdot (n - 2) = 2n - 2$ , completing the proof of Claim 1.  $\square$

Returning to our graph  $G$  with  $n$  edges and  $6n - 5$  edges, which satisfies (2), define  $G_1 = G\partial, G_2 = \partial G_1, G_3 = G_2\partial$ . Clearly, all the graphs  $G_1, G_2$ , and  $G_3$  satisfy (2) and hence, by applying Claim 1 three times, we conclude that the number of edges of  $G_3$  is at least  $6n - 5 - 3(2n - 2) = 1$ . Let  $e = [u_1, u_2]$  be an edge of  $G_3$ . Since  $G_3 = G_2\partial, [u_1, u_2]$  is a counterclockwise good edge of  $G_2$ . Consequently, there is an edge  $[u_1, v_1]$  of  $G_2$  such that the directed line  $\overrightarrow{u_1v_1}$  is obtained from  $\overrightarrow{u_1u_2}$  by rotating it counterclockwise around  $u_1$  by an angle smaller than  $\pi/100$  (see Fig. 1). Similarly, there is an edge  $[u_2, v_2]$  of  $G_2$  with  $\angle u_1u_2v_2 < \pi/100$ , as in Fig. 1. Since  $G_2 = \partial G_1$ , there are edges  $[v_1, w_1]$  and  $[v_2, w_2]$  of  $G_1$  with  $\angle u_1v_1w_1 < \pi/100$  and  $\angle u_2v_2w_2 < \pi/100$ , as in Fig. 1. (It is worth noting that it may be, for example, that  $[v_1, w_1]$  intersects both  $[v_2, u_2]$  and  $[v_2, w_2]$ , or even that  $w_1 = v_2$ .) Finally, as  $G_1 = G\partial$  there are edges  $[w_1, x_1]$  and  $[w_2, x_2]$  of  $G$ , with  $\angle v_1w_1x_1 < \pi/100$  and  $\angle v_2w_2x_2 < \pi/100$ , as in Fig. 1. All seven edges  $[x_2, w_2], [w_2, v_2], [v_2, u_2], [u_2, u_1], [u_1, v_1], [v_1, w_1]$ , and  $[w_1, x_1]$ , depicted in Fig. 1, belong to  $G$ . To complete the proof we show that they must contain three pairwise disjoint edges. Without loss of generality we may assume that  $\angle u_2u_1v_1 \geq \angle u_1u_2v_2$ .




Fig. 1

If the length  $l[v_2, u_2]$  of the segment  $[v_2, u_2]$  satisfies  $l[v_2, u_2] \geq l[u_1, u_2]$  (as is the case in Fig. 1), then we can easily check that  $[x_2, w_2]$ ,  $[v_2, u_2]$ , and  $[u_1, v_1]$  are three pairwise disjoint edges. Otherwise,  $l[v_2, u_2] < l[u_1, u_2]$  and then it is easy to check that  $[v_2, w_2]$ ,  $[u_1, u_2]$ , and  $[w_1, v_1]$  are three pairwise disjoint edges. Therefore, in any case,  $G$  contains three pairwise disjoint edges, completing the proof of Theorem 1.  $\square$

### Acknowledgment

We would like to thank Y. Kupitz, I. Krasikov, and M. A. Perles for helpful discussions.

### References

- [AA] J. Akiyama and N. Alon, Disjoint simplices and geometric hypergraphs, *Proc. 3rd New York Conference on Combinatorial Mathematics*, Annals of the New York Academy of Sciences, to appear.
- [AH] S. Avital and H. Hanani, Graphs, *Gilyonot Lematematika* 3(2) (1966), 2-8 (in Hebrew).
- [Er] P. Erdős, On sets of distances of  $n$  points, *Amer. Math. Monthly* 53 (1946), 248-250.
- [Ku] Y. S. Kupitz, *Extremal Problems in Combinatorial Geometry*, Aarhus University Lecture Notes Series, No. 53, Aarhus University, Denmark, 1979.
- [Pe] M. A. Perles, Unpublished notes.

Received May 19, 1988.